American. FRUIT GROWER

MAY

1946



for harve

uce foliage injupresent during th

same basic control long enough to ki ve been dusted.

YOUR WORK SPEEDS UP YOUR COSTS GO DOWN when you use FITESTOME *CENTER BITE* Ground Grips

YOU, as a farmer, know better than anyone else, the importance of keeping farm production costs down. You know just how important it is to do a seeding job or a harvesting job fast — when the time is right.

Firestone Ground Grip tires speed up work because they have more drawbar pull—up to 16% more. That's because they have no "open center" treads to pick up trash, clog up with mud and spin—dead in their tracks. They cut production costs because they have 40% longer tread life and a 14% stronger cord body.

Thousands upon thousands of farmers are doing more work at less cost today because they use Firestone Ground Grips. Ground Grips will do the same for you on every job you have, day after day, year in and year out. Get Ground Grips and prove it to your own satisfaction.

*Area in white shows the "Center Bite" traction zone, not found in other tires because of Firestone's exclusive patent rights.

Listen to the Voice of



Copyright, 1946, The Firestone Tire & Rubber Co.

DU PONT HELPS YOU b yourself to better crops

Control Insects

ers

lay

ps.

on

in

ve

ion

ne's

LONGER

CORD BODY

GROWER

DEENATE* DDT Insecticides. Your choice of DDT in four different formulations made for control of insects on the farm-including special forms for spraying and dusting fruits and vegetables. For easy application-effective results-and long-lasting control, follow this guide:

For Crops-Livestock-Buildings:

DEENATE 50-W (wettable powder, 50% DDT) DEENATE 25-W (wettable powder, 25° DDT)

For Livestock-Buildings:

DEENATE 25-R (Emulsifiable oil concentrate, 25% DDT) DEENATE 10-X (ready-to-use dust, 101: DDT)

Check Disease FERMATE* Fungicide. Non-caustic, nonrusseting on apples and pears. For scab and cedar-apple rust; brown rot and leaf spot of cherries and other stone fruits. Effective on many vegetables, too. Spray or dust.

ZERLATE* Fungicide. For brown rot of peaches; also early blight of potatoes and diseases of other vegetables. Spray or dust.

COPPER: A Compound. Effective copper fungicide, low in solubility for leafspot and brown rot of sour cherries; also many vegetables. Spray or dust.



DU PONT 2,4-D WEED KILLER. Kills many common weeds in pastures and lawns without harm to grass. Gets roots and all. Easy to mix-easy to apply as coarse spray-easy to wash out of equipment. New-different-effective.

U. S. Pat. Nos. 2,322,761 and 2,326,471

AMMATE* Weed Killer. For quick, permanent kill of poison ivy, choke cherry, wild blackberry, sumac and many other woody perennials.

Send for FREE Information on OU PONT PEST CONTROL PRODUCTS

E. I. du Pont de Nemours & Co. (Inc.) Grasselli Chemicals Dept., AF-4, Wilmington 98, Delaware.

Send information about the Du Pont Pest Control Products checked:

- **DEENATE DDT*** Insecticide
- **KRENITE*** Dinitro Spray
- ZERLATE* Fungicide
- COPPER-A Compound
- FERMATE* Fungicide 2.4-D WEED KILLER
- AMMATE* Weed Killer
- PARMONE* Fruit Drop Inhibitor

OTHER INSECTICIDES

- GRASSELLI* Lead Arsenate
- NUREXFORM* Lead Arsenate
- PARAPONT*
- Basic Lead Arsenate
- Calcium Arsenate
- Oil Sprays
- Paris Green
- ☐ ALORCO** Cryolite
- ☐ BLACK LEAF† "40" and "155"

OTHER FUNGICIDES

- ☐ SULFORON*
- SULFORON X*
- Lime Sulfur
- Flotation Sulfur Paste
- Copper Sulfate
- □ Bordeaux Mixture

OTHER MATERIALS

- Special Dust Mixtures
- Du Pont Spreader-Sticker
- Phenothiazine Worm Remedy
 - for Farm Livestock

Name

R.F.D. No. or Street.

Post Office_

State

*Reg. Trade Mark of E. I. du Pont de Nemeurs & Co. (inc.)

**Reg. Trade Mark of Aluminum Ore Company

+Reg. Trade Mark of Tobacco By-Products & Chemical Corp., Inc.

BETTER THINGS FOR BETTER LIVING ... THROUGH CHEMISTRY

DU PONT PEST CONTROL PRODUCTS

FAVORED FOR EVERY FARM JOB

Everywhere West
IN THE Mid-West

Farming's a business. And, like any other business, farming takes proper equipment, if a man expects to make money at it. On hundreds of thousands of farms, Ford Trucks are handling all the hauling jobs, heavy and light, around the place and in and out of town—all the way in to the big markets with loads of stock, poultry and crops. That's using headwork to save backwork—plus saving time and money!

Why do so many farmers pick Fords and stick to Fords? Simply because, for a quarter of a century Ford Trucks have been standing up under the toughest farm service, at mighty low cost. They've proved their economy and reliability in all kinds of work in all kinds

of weather. They're ideal trucks for farm service.

VOL

Lette

Vatio

State I Lesson:

Editoria

The new Ford Trucks are the best in Ford history. They're being delivered all over the country, right now—half-ton pickups and stake body trucks, tonners, heavy duty units—with 100-H.P. V-8 or 90-H.P. 6-cylinder engines—trucks with the newest engineering advancements.

Why not drop in at your Ford Dealer's next time you're in town, and see what Ford engineering has accomplished?

More Economical • More Reliable • More Enduring!

Check the exclusive features in the new Ford Trucks! They've got new fuel-and-oil-saving 4-ring aluminum alloy pistons, new longer-lived, steel-cored Silvaloy rod bearings, new cooling system protection to prevent loss of water and anti-freeze, new conveniences in servicing, new balanced carburetion to add to economy, new moisture-protected ignition—and a score of other important advancements. See the new Ford Trucks and get YOUR order in!

FORD TRUCKS



MORE FORD TRUCKS ON THE ROAD . ON MORE JOBS . FOR MORE GOOD REASONS

AMERICAN FRUIT GROWER

service. history. y, right trucks. V-8 or newest xt time ing has

MAY

VOL. 66

No. 5

during! Trucks!

ıminum Silvalov tion to convento add n-and

ASONS GROWER

CONTENTS Cover illustration courtesy U. S. Department of Agriculture Letters to the Editor The Great Mystery of Pollination 9 By Philip Ferry Problems of Orchard Pollination 10 By John R. Magness Looking Forward to the 1947 11 Apple Crop 12 Nationwide Fruits Peaches, Berries, Apples, Citrus, Nuts Hand Pollination of Delicious In the Wenatchee, Washington Orchards 14 By B. Esther Struckmeyer See the How A Beekeeper Sees It 16 By R. H. Kelty 19 American Pomological Society Founding of the American Pemplogical Society By R. L. McMunn 20 State News 20 Lessons in Orchard Chemistry Elemental Sulfurs Queen Shenandoah to Rule Blossom Festival 21 Ramblings of a Horticulturist 36 38 Editorial Page AMERICAN FRUIT GROWER Published Monthly by AMERICAN FRUIT GROWER PUBLISHING CO. 1370 Ontario St., Cleveland 13, Ohio E. G. K. MEISTER Publisher J. H. GOURLEY R. T. MEISTER ELDON S. BANTA (on leave) D. L. BARRETT Advertising Manager BRANCH OFFICES AND REPRESENTATIVES W YORK CITY, Billingslea & Ficke, 420 Lexington Ave. Room 506. Phone—Lexington 2-3667 AGO, Peck and Billingslea, 123 W. Madison St. Phone—Central 0465 SUBSCRIPTION RATES



ORTHO Scientific Pest Control is not merely a matter of materials. It is more than that. It is a year-round personalized comradeship of grower, dealer and ORTHO fieldman in the battle against horticultural diseases and pests. It is a scientific control of controls, varying with the individual need of the orchard or field.

for instance-DDT

This insecticide will undoubtedly have an important place in horticultural spray programs.

As yet, practical experience in general use is limited. Until all the implications of DDT are better known, growers will be well advised to proceed cautiously.

ORTHO field laboratories have tested and approved two formulations of DDT insecticides for 1946

RSISTO WETTABLE

PERSISTO DUST

ORTHO fieldmen can give you valuable assistance in introducing DDT into your spraying and dusting

RICHMOND



ELIZABETH

CALIFORNIA SPRAY-CHEMICAL CORPORATION



You can haul at a stepped-up fifth speed with McCormick-Deering Hauling Orchard tractors. The special five-forward speed transmission makes possible road speeds up to 15 miles per hour-a feature of these unique tractors that gives you great flexibility of operation. Both the O-4 and O-6 McCormick-Deering Orchard tractors are available with either high-compression, straight gasoline or distillate gasoline engines.

laneuverabil

Differential steering brakes and an extremely short turning-radius give these orchard tractors a high degree of maneuverability. They can slip in and out among the trees, around sharp turns between the rows. The differential brakes not only "snap" the tractor around quickly when necessary but they hold tight when you leave it standing on a hill or incline, Simple, easy-to-reach controls greatly reduce operational fatigue.

ultivation close to the trees

Smooth, streamlined surfaces prevent damage to young trees and trees with low-hanging branches even when you're working close to the trees. Branches and fruit glide right over the tractor without injury. Add the advantage of low overall height and you have a tractor ideally suited to orchard work.

Your International Harvester dealer will be glad to demonstrate these "extra" features of McCormick-Deering Orchard tractors. Let him show you how they will streamline your orchard.

INTERNATIONAL HARVESTER COMPANY 180 North Michigan Avenue Chicago 1, Illinois



McCormick-Deering Orchard

AMERICAN FRUIT GROWER

Dear Has

Dear F I ha

getting control just lik the lear bottom. the leav The produce

This atop the bottom or down the gun trees fr ous tops get it. Jackson.

Your facilitate sbrav n fair deg

Sawdu Gentlem Even magazin

magazin awmill, ground. sawdust this is a material

If the use of 1 happens. ing, now Hutchins

MAY.

ETTERS TO THE EDITOR

Wind As a Pollinator

Dear Sirs:

Has the wind a tendency to aid in pollination? Of course, the wet season of last year was a bad drawback, I know.

Scranton, Pennsylvania A Subscriber

The wind may be a factor in the crosspollination of fruit plants but it is usually of minor importance. Insects, especially honey bees, are the most active agents affecting pollination.

A wet, cool period during bloom often reduces the set of fruit because insects are prevented from making their normal visits to the flowers. Consequently there is a lack of good pollination.—Editor.

Controlling Apple Scab

Dear Editor:

I have read with much interest your article "Machinery for Applying Scab Control Sprays and Dusts." (Feb. '46.) It's getting closer to my experience but still off somewhat—like the man who didn't quite get into heaven. He ended up near-to-close, but just outside.

I have sprayed for many years from a 9 ft tower atop my spray tank. Had good control of scab from that level down but very poor indeed where I had to direct the spray upward. Scab spores floating through the air seem subject to gravity just like any dust so it is the top side of the leaves that gets infected and not the

through the air seem subject to gravity just like any dust so it is the top side of the leaves that gets infected and not the bottom. Examination shows that where the spray is directed upward, the tops of the leaves are very poorly covered.

The vigorous tops of my trees have not produced nearly so well as the lower weak

This winter I built a telescopic mast atop the spray tank. It is swiveled at the bottom and turned by a stick at the bottom. The same stick turns the spray up or down so everything can be reached. Thus the gun can be elevated to spray the tall trees from above or lowered to get the small ones. I want more from those vigorous tops and I think this year maybe I can get it.

Jackson, Michigan L. B. Benton

Your idea of using a telescopic mast to facilitate a more thorough application of stray material in the tops of the trees sounds very practical. Such homemade mechanical devices often work out to a fair degree of satisfaction.—Ed.

Sawdust As Mulch

Gentlemen:

Even though I raise fruit and grapes as a hobby in a very small way, I enjoy your magazine more than any other of some fifteen trade, sporting and other types of magazines I subscribe to.

As it happens, I also operate a small sawmill, cutting cottonwood. As you may know, this wood, in contact with the ground, is very perishable. Carloads of sawdust accumulate quickly. Why, since this is a non-resinous wood, would not this material be ideal for a mulch for orchard and vineyard?

If there are no facts available as to the use of this material, I thought I would mulch half of each planting and see what happens. Of course, I'd hate to do anything that would injure my little planting, now 4 years old.

Hutchinson, Kansas

E. H. Schroeder

Sawdust should make an excellent mulch for your small planting. Since it is largely a cellulose material and decays rather rapidly in the soil, it should serve equally as well as straw which is also a cellulose material.—Ed.

Pricing a Tree

Dear Sir:

I set out my prize tree orchard five years ago, using the best trees obtainable. Many are the exclusive Stark and Burbank strains; many are newly introduced and patented strains in the best varieties. From only two of my trees, when they were three years old, I sold twenty dollars worth of fruit. I have given my trees my best care and I prize them highly.

of fruit. I have given my trees my best care and I prize them highly.

My orchard is alongside of a main highway where I can sell all my fruit from my market at a nice profit. Among my trees I had a nice new Gorham pear, 5 years old, fruit as large as a cup, and the tree about 12 feet high and 8 feet wide

at the top.

A party drove off the highway and into my Gorham pear tree breaking it off at the base. In asking payment for the damage, I've valued the tree at fifty dollars (\$50.00). I'd like to have the opinion of other fruit growers if they think I'm right or if they think I'm too high. I'd like to know what you think the value of such a tree might be.

Some of you readers who have some real nice 5-year-old trees, just let me know at what value you believe yours to be worth either commercially or to your property.

Star Orchard

John L. Widner

Jacksonville, Missouri

What about it readers? What price would you put on your five-year-old trees?—Ed.

Salem Versus Danvers

Dear Sir:

I wish to call attention to an error in your geography in your last issue. (March

You say the "Famous Endicott Pear Tree in Salem, Massachusetts." It definitely is not in Salem. It is in the Danversport section of Danvers, Massachusetts. The tree is on the high ground above tide level on the North side of the Waters River, which at that point divides Peabody and Danvers. I learned to swim, did my first fishing and hunting of shore birds, trapped muskrats—in fact spent a whole lot of my kid's play there on the Peabody side of the Waters River, directly across the river fro mthe Endicott Pear Tree.

I'm rather hazy on the history now, having left that region 30 years ago, but I'm

I'm rather hazy on the history now, having left that region 30 years ago, but I'm quite sure that at one time Peabody was a part of Danvers. I almost think that Danvers was at one time a part of Salem. I'm not sure, though, of course, that could easily be substantiated. If that were correct the tree might originally have been in Salem, but to be correct now it would be necessary to say it is in Danvers:

Henniker, New Hampshire R. C. Coombs

The Endicott Pear Tree is located on Endicott St., Danvers, Mass. Endicott Street is in the section of Danvers, Mass., known as Danversport. In 1692, the city of Danvers was part of Salem. Mass. For this reason, the Endicott Pear Tree is often historically located in Salem.—Ed.



with the HANDIPILER

THE HANDIPILER loads, unloads, elevates, stacks, handles cartons, cases, bags, other packages. Compact, lightweight, safe, easy to operate, flexible, it is easily wheeled into crowded corners, around the shipping platform—fine for truck loading. Adjustable boom is clear of supporting obstructions; easily extended over piles or into car or truck. Floor locks hold machine in position. Reversible carrying apron. Available in four sizes, high end adjustable up to $7\frac{1}{2}$, $8\frac{1}{2}$, $9\frac{1}{2}$, and $10\frac{1}{2}$ feet; stacks commodities as high as 12, 13, or 14 feet. Handles individual items up to 100 lbs. Motor mechanism in base frame—plug into any convenient outlet. Write for Bulletin No. AFG-56.

STANDARD CONVEYOR CO.

General Offices: North St. Paul 9, Minn.





MAY, 1946

Deering

n makes

at gives

tractors

ctremely

ard trac-

around

around

incline.

hanging

over the

r ideally

tures of

gines.



COVER 'EM COMPLETELY

THE

connec

doubte

lination

continu

goes of

as we

nevert

living

ess of

most

pollen-

plants. broken

-food

that al

on the

part in

It is untold

have u

ing eac

al assis

plant.

the pol parts of prelimi

culmin

and fe

process transfe flower

mergin

MAY,

Polle

Commercial growers do not dispute the fact that the real profit on a crop is in a high "pack-out" of top grades. Too often, however, many growers lose sight of the vital requirement of timely and thorough spraying... and they lose the "money fruit" because of a "too little and too late" campaign against worms and scab.

Make this season's operation turn a better profit by keeping a step ahead of insects and diseases . . . by using the right spray materials, by putting them on at the right time, and, what is equally important, by using a sufficient amount of spray to completely cover the trees. For "Plus" Protection in your spray program, use dependable Orchard Brand products.

SPRAYS & DUST

Prevent Heavy Scab Losses...

MICRO-PARTICLE

ADDI C DDITOMIC* CIII EIII

APPLE DRITOMIC* SULFUR
the patented sodium thiosulfate feature that puts an "

With the patented sodium thiosulfate feature that puts an "extra wallop" into scab sprays. Built especially for apples and pears. Apple Dritomic Sulfur is unsurpassed for particle fineness, and is made to wet and disperse readily in the spray tank and to give high sulfur deposit on fruit and foliage.

Stop Codling Moth...

ORCHARD' BRAND LEAD ARSENATE

With the original fine, flake-like particle that results in better, more uniform deposit . . . increased protection against worm entries and stings.

GENITOX 550-50% DDT Spray Powder


Formulated after extensive laboratory and field studies augmented by General Chemical Company's background of nearly fifty years of insecticide manufacturing experience. Genitox S50 is of microparticle fineness, wets and disperses well in the spray tank, and gives a fine-divided "floc" resulting in maximum deposit on fruit and foliage and minimum run-off of the DDT insecticide. For use wherever DDT sprays are recommended or approved by State or Federal authorities.

GENERAL CHEMICAL COMPANY

40 RECTOR STREET, NEW YORK 6, N. Y.

Sales and Technical Service Offices: Atlanta - Baltimore - Boston - Bridgeport (Conn.) - Buffale Charlotte (N. C.) - Chicago - Cleveland - Denver - Detroit - Houston - Kansas Citt Los Angeles - Minneapolis - New York - Philadelphia - Pittsburgh - Providence (R. L) San Francisco - Seattle - St. Louis - Utica (N. Y.) - Wenatchee & Yakima (Wash.) In Wisconsin: General Chemical Wisconsin Corporation, Milwauken. (Wash.) In Canada: The Nichols Chemical Company, Limited - Montreal - Toronto - Vancouver

*Reg. U. S. Pat. Off. †Trade Mark. General Chemical Company



By PHILIP FERRY

Honeybees at work in a cluster of blossoms.

1

uts an "extra

es and pears. neness, and is

and to give

ENATE

n better, more m entries and

Powder -

es augmented ely fifty years do is of microay tank, and bosit on fruit cide. For use d by State of

PANY

(Conp.) • Buffalo on • Kansas Chy Providence (R. I.) kima (Wash.) e, Wis.

GROWER

HE most mysterious phenomenon connected with the plant world is undoubtedly that of pollination. Pollination is a dramatic, a vital and a continuously operating process which goes on all through nature. Unaware as we may be of this phenomenon, it nevertheless continues unabated. All living creatures must eat; in the process of procuring food, all insects and most animals aid in disseminating pollen-the latter the lifeblood of plants. The whole operation is an unbroken cycle: animal—plant—pollen—food. Investigation has revealed that almost every creature that moves on the face of the earth plays some part in the perpetuation of plant life. It is interesting to reflect that for untold centuries plants and insects have unknowingly been complementing each other in this process of mutual assistance.

Pollen is the male product of the plant. Pollination is the transfer of the pollen from the male to the female parts of the flower and is merely a preliminary to other changes which culminate in fertilization. Pollination and fertilization are very different processes. Pollination is merely the transfer of the pollen grains from flower to flower; fertilization is the merging of the male element or sperm

and the egg nucleus. The union of these germ cells brings about the fertilization of the eggs in the ovules of the ovary and the subsequent development into seed.

There are two distinct types of pollination: self-pollination (autogamy), in which pollen is transferred from the stamen to the pistil of the same flower: and cross-fertilization (allogamy) in which the pollen is carried to the pistil of another flower on the same plant or to another plant of the same species. As the pollen-bearing stamen is rarely in contact with the female stigma at the time when both of these are ripe, some means is clearly needed to bring the pollen to the female organ. There are countless methods of accomplishing this union, the more common being pollination by insect, wind, bird, or animal agency, and even pollination by water. Perhaps the most celebrated carrier of pollen is the bee; the most picturesque the hummingbird; the most spectacular the Pronuba moth which fertilizes the Yucca plant of western deserts.

Even the lowly ant, slug and snail participate in this universal phenomenon. Squirrels, by forgetting where they hide part of their trove also aid, their poor memory being an important factor in the propagation of forests.

to a person's garments, that person automatically becomes a disseminating agent; although the seed pods were

designed to cling to the hooves and hides of animals, and although man was not considered in the original design, such a person assists wily nature in the scattering of seed and pollen—

a highly important function in nature. Insects are the chief agent in the work of pollination, and the great variety in color, form and scent of flowers has been developed in relation to the precise insect pollinator which serves each particular species. When we speak of the great mystery of life, we must include insects as one of the chief instruments in the furtherance of this unfathomable design. Most plants pollinated by insects have conspicuous flowers or a distinct odor or both. All such flowers have glands which secrete nectar, the latter the lure that brings insects to plants. Nectar is the sugary liquid from which bees manufacture honey, and there are more than 2000 different plants growing in the United States from which the insects extract the substance. The term is derived from the name given by the Greek poets to the drink of the ancient gods. Without the plant and its nectar, the in-

(Continued on page 31)



indicated in the table. Generally early and medium-blooming varieties will cross pollinate satisfactorily, and medium and late, but the over-lap of bloom between early and late bloomers may not be sufficient for satisfactory pollination.

Many orchards now in production do not have suitable pollination conditions. Large blocks of Winesap and Stayman, or solid Delicious or Mc-

Left—Pollinating "bouquet" method on low tripod in a solid block of York Imperial apples. Photo by Bureau of Plant Industry, Soils, and Agricultural Engineering, U.S.D.

char

Of the or grow weat part

befo rain

with loss fruit

drop

char

mean

seas

prop

trien

and

only

by tl

to de

sibili

befo

This

buds

live

thro

twig

It

PROBLEMS OF ORCHARD POLLINATION

By John R. Magness, Head Horticulturist in Charge, Division of Fruit and Vegetable Crops and Diseases, Bureau of Plant Industry, Soils, and Agricultural Engineering, Agricultural Research Administration, U. S. Department of Agriculture

ABOUT half a century ago, it was first discovered that we have a pollination problem with certain of our orchard crops. At that time, Mr. M. B. Waite discovered that a number of pear varieties would not set fruit unless fertilized with pollen from another variety. His work introduced the pollination problem into orchard management. Prior to that time orchards generally included many varieties, and the pollination problem was not recognized.

Since Dr. Waite's early work, pollination studies have been made on all important fruit crops. We now know that some kinds of fruits are generally self-fruitful, setting full crops with their own pollen. These include the citrus fruits, peaches, apricots, sour cherries and most of the Domestica or European plum and prune varieties.


We also know that other kinds generally are not fully self-fruitful, and cross pollination is essential to a full set of fruit. This group includes apples, pears, sweet cherries, and Japanese plums. Within this group, some varieties will set partial crops without cross pollination but for full production, crossing is needed. We also have some varieties, particularly in sweet cherries and Japanese plums, that are inter-sterile. Thus with these kinds of fruits provision for cross pollination should always be made when the orchard is set.

The table on this page lists the 24

leading apple varieties of the United States, with the season of bloom, and whether or not they are good pollen producing varieties.

In planting the orchard, varieties should be so arranged that all trees are within 100 feet or less of a good pollinizer. Thus not more than four rows of a variety should be planted in a block, with suitable pollinizers on each side. Varieties listed as poor pollen producers should not be depended on for pollination purposes.

The general season of bloom is also



Intosh orchards frequently do not set full crops for this reason. What steps can the grower take when he has such a condition?

First of all, if the orchard is not too old, he should take steps to permanently correct the problem by grafting over certain trees to good pollinizer varieties. Every third tree in every third row is a safe recom-

(Continued on page 28)

Variety	Season of Bloom	Value as Pollinizers
Delicious (and color strains)	Medium	Good
Winesap " " "		Worthless
Jonathan """		Good
Stayman Winesap (and color strains)		Poor
Ben Davis (and color strains)	39	Good
D D . # # #	Late	Good
7 1 T - 1 1 01 42 40	Medium to late	Good "
AF.T., 1. NO AN AN	34 - 41	
Grimes Golden		**
Yellow Newtown		**
Wealthy		**
Yellow Transparent		89
Rhode Island Greening		Poor
Northern Spy		Good
Gravenstein		Poor
Duchess of Oldenburg		Good
Arkansas (Black Twig)	Medium	Poor
Golden Delicious	"	Good
Esopus Spitzenburg	**	"
Vagener		"
btark	34 11	Poor
Vinter Banana		Good
Cortland	29	"

e. Generally ning varieties factorily, and the over-lap rly and late sufficient for

in production llination con-Winesap and cious or Mc-

York Imperial Plant Industry, eering, U.S.D.A.

pollination by in his trees.

ntly do not ason. What ke when he

hard is not the steps to problem by the to good by third tree to afe recom-

lue as linizers

ood rthless ood oor ood ood

oor ood oor

oor ood ""

,, ,,

GROWER

LOOKING FORWARD
To The 1947 Apple Crop

By A. J. HEINICKE New York Experiment Station

THE title of this article implies that it may not be too early for the orchardist to think about the 1947 crop. Of course, we are not yet sure what the crop will be in 1946. Every fruit grower knows that unfavorable weather from now on until the latter part of the growing season can blast the prospects for a good harvest.

Such hazards as late spring frosts, before or after buds open, cold or rainy weather during bloom that injures the flower parts or interferes with pollination, a heavy "june drop," loss of a large part of the growing fruit from insects and diseases, from hail and wind, or from a premature drop just before harvest are all too familiar experiences in many orchards. A good set of clean fruit in early summer does not necessarily mean a heavy crop, especially in dry seasons. If the apples fail to size properly due to lack of water or nutrient materials supplied from the soil and by the leaves, the crop may be only 1/2 to 2/3 as large as indicated by the set in early summer.

It is a very simple matter, however, to determine whether or not the possibility for a crop exists many months before the flowers normally open. This can be done by examining the buds to see whether or not they have live flowers. If present, the flower parts can be exposed by cutting through the buds while still dormant or after they begin to swell. A few twigs may be removed from the trees



Right—Bees are very effective agents in carrying pollen from tree to tree during bloom. Overpollination is to be avoided whenever possible.


and placed with the cut ends in a jar of water and forced in a warm room. If there are any flowers in the buds, they will appear within a few weeks' time. If there are no flowers present in the buds, it is impossible to have any fruits in 1946.


Most fruit growers know that the flower buds that open in spring begin to form very early in the previous season. Through microscopic examination of the very small buds, it has been shown that the embryonic parts of the flowers may be present as early as the latter part of June, approximately 8 or 9 months ahead of the time that they open. Not all the flower buds, however, begin their develop-




ment at the same time, and initial stages may be observed for several months, in some cases as late as August. Once a flower bud has been started on its way, it may undergo (Continued on page 22)

Below left—Fruit spurs from an apple tree of low vigor. None of these spurs show indications of blossoming. Below right—Fruit spurs from a vigorous tree of the same variety. These spurs are making a satisfactory growth, and blossom buds are unfolding.







Pear blossom scene in Hood River Valley, Oregon, with Mt. Adams in the distance.

PEACHES

Heavy Pruning Lowers Production

As recently reported in the Missouri "Horticultural News" by Dr. Aubrev D. Hibbard of the University of Missouri's Horticulture Department, studies in progress at the Campbell Field on pruning peach trees show that the removal of even a moderate amount of wood may materially lower production. In his zeal to properly train the young trees, the grower may be literally cutting away his future crops, says Dr. Hibbard. The undesirability of heavy pruning of young trees has been indicated by pruning experiments starting with three year trees and continuing until the trees were eight years of age. Three degrees of pruning were given to the trees. The heaviest was called "moderate" which is the heading and thinning type usually recommended. Trees that were thinned but not headed were designated as "light" pruning. Only the interfering, dead, and broken branches were removed in the third kind of pruning which was labeled "corrective."

As determined by the weight of

the wood removed, the light pruning was twice as severe as the corrective type while the moderate was three times as severe. Dr. Hibbard reports that before pruning in 1945 the trees in all treatments were of approximately the same height despite annual heading of the moderately pruned trees. Trunk measurements revealed that the trees given the lightest pruning were slightly over one-half inch' larger in circumference. With respect to cultivation, fertilization, spraying, and thinning, the trees have been given identical culture.

Although four crops have been harvested from these trees, there has been no difference in the quality of fruit under any treatment. Total yields per tree for the four seasons have been 17.2 bu. for the corrective pruning, 16.4 bu. for the light, and 14.7 bu. for the moderate. These differences seem small, says Dr. Hibbard, but when the loss per acre is calculated they assume considerable proportions. If the crop had been worth only \$1.50 per bushel on the tree, the loss from light pruning would be \$91.50 while that from moderate pruning would amount to \$285.00 per acre.

In conclusion, Dr. Hibbard reports that not only was there a loss of fruit but the moderately pruned trees required three times as much labor for pruning. So far there has been no greater breakage in the corrective pruned trees. Moreover the fruit has not been any more difficult to harvest.

BERRIES

Pruning Small Fruits

If small fruits are to yield well and be easily kept in bounds, all except the strawberry must be pruned regularly, says W. H. Childs, associate horticulturist at the Agricultural Experiment Station, West Virginia University, Morgantown, West Virginia. Raspberries and blackberries must be pruned during the summer as well as in early March.

In pruning black raspberries, the lateral branches should be cut back to about 8 to 12 inches, and any weak canes should be removed. If the plants were not topped last spring, the best a grower can do is to cut the canes back enough so that he can cultivate and harvest the fruit without undue trouble.

Purple raspberries are treated the

turple raspoerries are treated in

12

AMERICAN FRUIT GROWER

longer thinne in the ceed fi hack t in the sembl raspbe latera inches thinne in the Can be ren blackt been d Wit

eame

latera

any ca and lin old ca eight. moved Witt able to patch often s

Olde

prunir

In th ington of mo Its his Bancre of it: about young embar employ seeds slipped the wa men: destina haulin

garder Mrs cus W ing his 12, 18 planter severa a long the on! In the

died. le

Eve

seeds

seems forgot comma that si ground largely specto berg,

and id

same as blackcaps except that the laterals are left about two inches longer. Red raspberries should be thinned to about 6 or 8 inches apart in the hedge-row, and any that exceed five feet in height should be cut back to that height if they bend over

The pruning of blackberries resembles that of both red and black raspberries in some respects. The laterals should be cut back to 18 inches in length, and the plants thinned to about 6 to 8 inches apart

in the hedge-row.

Canes which bore last year should be removed from all raspberries and blackberries, if this has not already been done.

With currants and gooseberries, pruning consists chiefly of removing any canes more than three years old, and limiting the number of one-yearold canes from the crown to about eight. Any dead wood should be removed also.

With all these fruits, it is advisable to carry all prunings from the patch and burn them, since they often serve as a source of disease.

Oldest Apple Tree of Northwest

In the Vancouver Barracks, Washington, there stands an apple tree of more than ordinary importance. Its history was so interesting that Bancroft, the noted historian, says of it: "At a lunch party in London, about 1825, given in honor of some young gentlemen who were about to embark for Fort Vancouver in the employ of the Hudson Bay Company, seeds of the fruit eaten were slyly slipped by some young ladies into the waistcoat pockets of the young men. Upon their arrival at their destination, the young men, in overhauling their wardrobes, found the seeds and gave them to Bruce, the gardener at the Fort."

Mrs. Mary Whitman, wife of Marcus Whitman, also wrote an interesting history of this tree on September 12, 1836. It is said that the seeds planted by the gardener produced several trees, three of which lived for a long time, and were pointed out as the only apple trees in the Northwest. In the course of time two of them died, leaving the present tree alone.

Even the existence of this tree seems to have been almost entirely forgotten by the public, not even the commander of the barracks knowing that such a tree stood on the very ground which he controlled. It was largely through the horticultural inspector of this district, A. A. Quarnberg, that the tree was discovered and identified. In 1911, Mr. Quarn-

MAY, 1946



The sign in front of the tree reads: "The oldest apple tree in the Northwest grown from seeds brought from London, England, planted in 1825 by the Hudson Bay Co.

berg wished to have a gavel made of wood from the tree for the Washington State Horticultural Association, and, upon examining the tree, found it badly infested with San Jose scale, half of its branches dead, and in a bad condition generally. On January 13 of that year he called upon Col. G. K. McGunnigle, Commander of the Barracks, and got the necessary permit to prune, spray, and do anything necessary to preserve the life of the tree.


On January 25, 1911, by direction of the Washington State Commissioner of Agriculture, he took measurements and found the tree to have the following dimensions: one foot from the ground, diameter 11/2 feet; height 33 feet; spread of crown 33 feet. On February 21, 1911, the tree was pruned and all dead branches and brush removed, the decayed part of the trunk and branches cleaned out and filled with plaster of Paris and cement, and all cuts painted.

The tree is evidently a Yellow Bellflower seedling. In 1915 the Department of Agriculture at Wash-. ington, D.C., requested Mr. Quarnberg to send them specimens of apples for making wax forms. The picture on this page shows this tree after it was pruned .- H. E. Zimmer-

The Fruit With the Zipper Skin

The same Italian sailor who is credited with the discovery of the New World in 1492 also brought the seeds of citrus fruits from Spain to the Canary Islands on his second voyage, and years later the Spanish conquistadores carried the fruits to Florida where they have flourished

The tangerine has made a definite place for itself as a favorite winter fruit—the fruit with the zipper skin -in spite of the greater volume of its sister citrus, the orange and



bbard renere a loss ely pruned as much there has in the coreover the re difficult

its vield well ds, all exbe pruned is, associricultural Virginia West Virckberries summer

cut back and any ed. If the st spring. is to cut at he can uit with-

rries, the

eated the



HAND POLLINATION OF DELICIOUS

In The Wenatchee, Washington Orchards

By B. ESTHER STRUCKMEYER

University of Wisconsin

DIFFICULTIES have been experienced in the setting of fruit in Delicious growing in Wisconsin. There are some known peculiarities of Delicious and some possible ones which might contribute to the difficulty of getting this variety to set satisfactorily. One of these which contributes to the trouble is blossom structure. In the AMERICAN FRUIT GROWER for April, 1945, Dr. R. H. Roberts described the structure of the Delicious blossom. He showed that the structure of the blossom permitted bees to extract nectar from the base of the flower without having to crawl over the stigmas and stamens, resulting in · the failure of the bees to pollinate the blossom. The petals of this variety are flat and assume a position at approximately right angles to the upright pistil and stamens, so that it is not necessary for bees to enter the flower through the pistil and stamens in order to reach the nectaries. In apple varieties where the filaments bearing the anthers are wide spread or the petals are cupped, bees reach the nectaries only by passing over the stamens and pistils, thereby pollinating the flowers. Varieties with this type of blossom set fruit readily with bee pollination. Furthermore, the

pistils on the Delicious blossom are short, so that bees may not touch the stigmas even when collecting pollen.

A second peculiarity of Delicious is the observed preference bees may have for other varieties of apples or plants blossoming at the same time. Although there may be enough bees in the orchard for good pollination, they may be attracted to other varieties of apples in preference to Delicious to the extent that there is an insufficient number visiting Delicious trees to effect adequate pollination. It has been suggested that the nectar in Delicious is not as attractive as it is in other varieties, or possibly the nectar is desirable to bees for only a short time instead of the entire blossoming season. As Delicious is self-sterile, it is also important to have good pollinizers close to the Delicious trees.

Since bees are important in cross-pollination, why not have enough of them to overcome the handicaps to which Delicious may be subject? This would entail the expense of bringing in bees, but probably more important is the fact that bringing in an abundance of bees for Delicious is apt to result in over pollination of other varieties, and since over pollinated trees need hand thinning it is apparently

Left—Blossom time in the Cashmere Velley near Wenatchee, Washington.—Photo from the Liberty Orchards Co., Cashmere. (Above)—The central blossom in the clutter is usually pollinated shortly after the petals have unfolded. It is important that the pollen is placed on all five stigmas.



In collecting the pollen for hand pollination, a flower is removed from the spur and rubbed over a mesh screen so that the anthers fall into the collector's jar.

cheaper to hand pollinate and cut down thinning expense.

Hand pollination is a common practice in the Wenatchee Valley in Washington. On a visit to this section the writer was able to observe the collecting, curing and application of pollen to the blossoms. Some of the common varieties used as sources of pollen for Delicious in Washington are Jonathan, Golden Delicious, Newtown, Winter Banana and Rome Beauty.

Hand pollination in Washington is done on a commercial scale. The first step is collecting the pollen. The dealers who make a business of gath-

(Continued on page 32)

Healthy LEAVES Produce Superior Fruit



YOU WILL PICK BETTER FRUIT

if 30 or more healthy leaves nourish each apple.

Leaf injury by aphids and leaf hoppers reduces fruit yield and quality. Leaf injury from insects commences early in the season and leaf efficiency for the remainder of the season is considerably reduced. Black Leaf spray programs help to protect healthy leaves, by controlling aphids and leaf hoppers.

continue for nicotine
products essential in
We suggest that
their nicotine supply
tant protective sprays
of the growing
season.

Black Leaf 40 and Black Leaf 155

are not caustic or toxic to foliage or fruit, and leave no objectionable residues at harvest.

Black Leaf 155

Black Leaf 155 controls leaf hoppers and , leaf miners as well as codling moth . . . lets healthy leaves build yields and profits.

Experienced growers say its performance makes it profitable.

TOBACCO BY-PRODUCTS & CHEMICAL CORP. INCORPORATED · · LOUISVILLE 2, KENTUCKY



The first en. The s of gath-

nd polline-

and cut

mon pracin Wash-

ection the

of pollen the com-

es of pol-

ngton are

d Rome

nington is

ROWER



Four truckloads of honeybees, contracts for on a rental basis, moving into an or chard for distribution at pollination time

HOW A BEEKEEPER SEES IT

Beekeepers' Viewpoint on the Pollination Problem

By R. H. KELTY, Michigan State College

SINCE January 1, I have conducted beekeepers' meetings in approximately thirty Michigan counties, in and out of the Michigan fruit belt. The situation calls for frank appraisal and discussion. Beekeepers are apprehensive about spray poisoning. Many who have rented bees formerly will not do so this year because of their sorry experience last season. Many of those who are willing to try it again are demanding a higher rental price to offset increased labor costs and risk of poisoning. There is a strong ten-dency among larger beekeepers in the fruit belt to move their bees completely out of the commercial orchard area until more is known about the effect of DDT on bees in orchards.

In our opinion, fruit growers should prepare to handle their pollination problems in a more orderly manner. Random rental of a few hives of bees here and there, on the last days before the blooming period, without assurance of strength of colony, food supply or time of delivery and removal of the bees from the orchard, is a loose business arrangement.

The beekeeper who depends upon his bees for a living, depends upon the strength of his colonies for honey production. The beekeeper who keeps a few bees without much attention, may not even know whether his colonies are strong or weak or hungry or free

from disease. To secure a supply of bees in quantity for pollination, fruit growers may need to contact commercial beekeepers, most of whom are not located in the fruit belt. This may involve transporting the bees a considerable distance at added expense, and it may involve a higher rental price to cover actual costs, but it is likely that the fruit grower will get more for his money in actual pollination service if he deals with estab-

lished, successful commercial honey producers whose income depends upon strong colonies.

The fruit grower should expect to get strong colonies of bees, capable of heavy bee flight on days of favorable weather conditions during the blooming period. From such colonies. there may be from 100 to 200 bees flying per minute when the temperature is above 70, on bright, calm, balmy days. Under the same conditions, from weak colonies there may be flying less than 50 bees per minute. Strong colonies containing six or eight or more combs of brood, need a reserve food supply of twenty or more pounds of honey, or they may starve in mid-bloom in cool rainy weather. The bees should be properly prepared for moving so that inexpenenced helpers can handle the hives without danger of stinging. Comparatively sheltered spots should be chosen in the orchard for the bees, (Continued on page 33)



A bee yard on a Clinton County farm in Ohio.

NEW GMCs for Every Farm Use



and IMPROVED

Powerful, economical "Army Workhorse" engines of Valve-in-Head design with Tocco-hardened Crankshafts, Airplane type "Durex" Main Bearings, Heat-resisting Exhaust Valves with improved Valve Seat Inserts, Positive Crankcase Ventilation, Turbo-Top Pistons and Full Pressure Lubrication through Rifle Drilled Connecting Rods.

 Rugged, built-for-the-job chassis with extra strong Frames, Springs and Axles, heavy duty Transmissions and Clutches, Needle Bearing Universal Joints, Recirculating Ball-Bearing Steering and powerful, easy-action Brakes.

• Rider Ease Cabs with Cradle-Coil Cushions, Wide Visibility ""V" Windshields, Quick-vision Instrument Penels, All-Weather Insulation and Controlled Ventilation.

New GMCs offer you the most in farm truck value in a most complete selection of models . . . 1/2 to 20 tons. There are pick-ups in several sizes . . . all-steel with grain-tight gates. There are panels in capacities of 1/2 to 11/2 tons, with fully insulated sides and tops to protect perishables. There are stakes, platforms and tractor units in tonnage ranges and with equipment options to answer every other farm t.ansport need. Whatever you haul . . . grain, milk, poultry, fruit, vegetables, livestock, feed, fertilizer and many more...you'll find a war-proved, improved GMC ideally suited to the job.

THE TRUCK OF VALUE



GASOLINE

GMC TRUCK & COACH DIVISION . GENERAL MOTORS CORPORATION

MAY, 1946

17

ercial honey me depends ald expect to bees, capable lys of favorduring the such colonies. to 200 bees the temperbright, calm, same condies there may s per minute. ning six or rood, need a twenty or or they may cool rainy l be properly hat inexperile the hives

ging. Coms should be

or the bees, e 33)

ETTER PEST CONTROL

MICRONIZED







WRITE FOR FURTHER INFORMATION

SHERWIN-WILLIAMS SPRAY MATERIALS

INSECTICIDE DIVISION

101 Prospect Ave., N. W.

Cleveland, Ohio



THE ican I proach question to the tors of the societi To ap the Societi To ap the societi the ing in this with the tors of the societi the ing in this with the societi the ing in this with the ing in the ing

II and
of the
dated t
ical Co
take up
Societiologica
first fe
Althe
cultura
prior t

was, as

rules f varieties recting so great known men, g realized or noth it out a

Short "Fruit a was pul Humric had an Horticu which I ing's wrules that

fruits ar

MAY,



Founding of the American Pomological Society

PART I: RULES OF POMOLOGY

By R. L. McMUNN Department of Horticulture University of Illinois, Urbana, Illinois

HE Centennial year of the American Pomological Society is fast approaching. Because there is some question as to the correct date of the organization of the Society it seems advisable to trace the events and mention the factors or forces leading up to its organization. One of these factors was the chaotic condition of pomological nomenclature. Another was the previous existence of two societies which could be consolidated. To appreciate better the founding of the Society it seems advisable to present the facts in four parts, each part being presented in a different issue in this magazine. Part I will deal with the adoption of rules of pomology or nomenclature. This subject was, as it is now, of prime importance to nurserymen and pomologists. Parts II and III will sketch the early history of the two organizations that consolidated to form the American Pomological Congress (Society). Part IV will take up the consolidation of the two Societies to form the American Pomological Congress (Society) and the first few meetings of this Society.

Although there were several horticultural societies in the eastern states prior to 1846 none apparently had rules for their guidance in naming varieties, establishing priority, or correcting synonyms. The confusion was so great that a single variety was often known under many names. Nurserymen, growers and pomologists alike realized the great confusion, but little or nothing could be done to straighten it out as long as there were no generally accepted rules.

Shortly after A. J. Downing's book, "Fruit and Fruit Trees of America' was published in 1845, a. Mr. T. S. Humrickhouse of Coshocton, Ohio, had an article in the Magazine of Horticulture of February, 1846, in which he criticized parts of Downing's work, then proposed a set of rules that could be followed in naming fruits and establishing priority. This, o far as is known, is the earliest sug-

gestion of rules of pomology. In late September, 1847, a group of growers and nurserymen assembled at Columbus, Ohio, and organized the "Ohio Nurserymen and Fruit Growers' Convention." This organization adopted a set of pomological rules. Whether the set of rules was the same as that proposed by Humrickhouse is not known. Downing in the magazine Horticulturist of April, 1848, said the rules were the same as those proposed by the Massachusetts Society. (Downing was incorrect in this statement because the Ohio Convention adopted its rules before the Massachusetts Societv proposed its rules).

Whether the rules suggested by Humrickhouse prompted Downing to propose a set of rules is not known. At any rate, in the October, 1847, issue of the Horticulturist, Downing ran an article entitled "Pomological Reform" in which he suggested a number of rules that might be adopted by all societies for the sake of uni-formity. In December, 1847, Downing had the pleasure of announcing in



Andrew Jackson Downing, famous American pomologist and landscape gardener.— Photo Massachusetts Horticultural Society.

the columns of the Horticulturist that the Massachusetts, Pennsylvania and Cincinnati Societies had adopted a series of "Rules for American Pomology." This set of rules was soon adopted by several other societies. It is this set of rules that the American Pomological Society later adopted. In January, 1846, the New York

State Agricultural Society appointed a fruit committee to bring in a list of recommended fruits. This committee reported the following year and then requested that it might continue its work for another year. This request was granted. At a committee meeting of the Agricultural Society on March 11, 1847, Downing's "Fruit and Fruit Trees of America" was made the authority of the Agricultural Society. Also at this meeting, Downing and J. W. Bissell were named to the fruit committee. During the Agricultural Society committee meeting on January 21, 1848, the fruit committee reported a "Set of Rules of Pomology' which were very short and varied somewhat from the "Rules for American Pomology," the set of rules that nearly approached the suggestions made by Downing in October, 1847. A few societies soon adopted the New York Society rules. Also at this meeting the fruit committee reported, -and they also regret the nonreception of the lists recommended by those gentlemen (i.e. Downing and Bissell) until after the lists appended hereto was adopted, otherwise some additions might probably have been made to it,-." It seems probable that the "non-reception" of the Downing-Bissell list of fruits and the fact that the other members of the fruit committee presented a set of rules that were different from those suggested by Downing at an earlier date may have been the incentive for the animosity Downing showed towards the Agricultural Society, the Society that sponsored the North American Pomological Convention, one of the parent organizations of the American Pomological Congress (Society).

Much controversial material was written about the relative merits of the different rules. C. M. Hovey, editor of the Magazine of Horticulture was champion of the New York rules; Downing was the standard bearer for the Massachusetts rules. This was the start of the ill feeling that was to last for two years. It was probably one of the reasons why two national conventions organized in the fall of 1848, rather than meeting as a single or-

Regardless of the merits of the different rules, the credit for proposing the first set of rules must go to Mr. T. S. Humrickhouse, and the laurels for adopting the first set of rules to the Ohio Convention.

NEWS state

MAINE—Another Farm and Home Week at the University has been held. The Pomological program dealt with economic trends, tree vigor, apple insect control, and apple processing. Dr. J. K. Shaw of Massachusetts and Dr. J. J. Willaman of the Eastern Regional Research Laboratory in Pennsylvania spoke on this program. In addition, Dr. Willaman visited the chemical engineering laboratories where some pro-cessing is underway, and addressed an eve-ning meeting on "The Work of the Regional Laboratories in Finding New Uses for Farm Products."

Kinds and varieties of fruits suitable for home freezing, and means of making a success with fruits in the home garden were scheduled in programs other than pomo-

Dr. Joseph A. Chucka, Agronomist and head of the department of soils and crops in the experiment station, has resigned and moved to Springfield, Mass., where he is associated with Eastern States Farmers' Exchange. Dr. Chucka had, in recent years, been devoting some attention to problems of orchard fertility and made a very real contribution in demonstrating that a leaf disorder long recognized in this State and called *leaf scorch* really results from a deficiency of magnesium, and that the symptoms disappear and the trees again become healthy when treated with magnesium in the form of Epsom salts. We share his confidence that applications of magnesium limestone will do the same, but more slowly.—J. H. Waring, Professor of Horticulture, University of Maine, Orono.

MICHIGAN—The outlook for fruit in Michigan for 1946 is very good. Moderate winter temperatures have been favorable. Peach trees have come through the winter with very little injury to the buds. Barring a repetition of last year's spring weather, the fruitfulness of Michigan orchards should be above average. Up to the twentieth of March the season has been a repetition of 1945 and there is considerable concern over the possibility of an early bloom.

The interest shown by the attendance of 1600 fruit growers at four district Horticulture Society meetings held in the State this spring is some indication of the desire by the growers to know the new develop-ments in the fruit business of Michigan and

other fruit producing states.

A very keen interest is shown in the new insecticide, DDT. The indications are that a large number of growers will adopt the DDT spray schedule for apples, pears, peaches and grapes. Experimental work with DDT in the State in 1945 was done with 25% wettable powder, 3 pounds per hundred gallons of water, which gave satisfactory control of codling moth, leaf hop-per, Oriental peach moth, grape berry moth, rose chafers, and tarnished plant bugs. Some control was also achieved on green apple aphis and black aphis on sour cherries.

Apple scab took its toll of foliage on many apple orchards in 1945. Some of the most badly defoliated orchards will probably not blossom this year. Other orchards will be in varying stages of productivity, depending upon their state of vigor, but in general the outlook is good as of March 20.

—C. A. Langer, Extension Specialist in Horticulture, Michigan State College, East Lansing.

NEW YORK—As reported in a recent issue of the New York State Horticultural News Letter, edited by T. E. LaMont of Albion, Colin Nichols of Lewiston was elected president at the annual meeting of the New York Cherry Growers' Associa-tion. Other officers elected were: Earl Harding of Albion, vice-president, and Snow Wooster of Ontario, secretarytreasurer.

Dues for the year 1945 amounted to \$2,814 and the membership totaled 898. One hundred and forty-one members came in from the newly organized district around

Erie, Pennsylvania.

The plan of cooperating with the New York State Horticultural Society in employing a full time executive secretary for the fruit industry of the State was approved by the Cherry Growers.

PENNSYLVANIA - Preliminary plans for the construction of a new laboratory building featured the annual meeting of the advisory committee of the Arendtsville Fruit Research Laboratory held at Peach Glen recently. The advisory committee, consisting of representatives of the fruit growers associations in Adams, Franklin, York and Cumberland counties, counsels with the workers from the Pennsylvania State College Agricultural Experiment Station regarding research conducted in the fruit region of the state.

The new laboratory building, according to present plans will cost about \$25,000 and will be financed by funds contributed on a non-profit agreement by individual fruit growers. The experiment station plans to rent the building from the growers, F. F. Lininger, Director, explained, with the understanding that the College may accept title of ownership at some later date.

OHIO—Recognizing the service of the Ohio Pomological Society, which served Ohio fruit growers from 1847 to 1867 when its officers and members continued on as the Ohio State Horticultural Society in order to broaden the base of operation and serve more horticultural interests, the Ohio State Horticultural Society decided at their annual meeting in Cincinnati to recognize 99 years of service to the Ohio fruit industry and prepared to celebrate 100 years of service with a Centennial Meeting next winter. A Centennial Meeting of the Ohio State Horticultural Society is scheduled for February 25, 26 and 27, 1947, at the Deshler-Wallick Hotel, Columbus, Ohio.

Thomas E. Thornburg, Ashland, Ohio, oldest living member of the Society was

OUL

BL

The

Rlosson

chester,

will hav

Miss N

Secreta

Clinton

first qu

son wil

000 loy

to gathe

day cele

whose I Mexico Semina

active glee clu

also se

presider

worker

taken c

with th

the sixt

ton's a

tacting i

annually

Festival

coronati

Court,

housing

queens

Matthey

Miss Pr

chester. ington;

Belgium

Miss N

vakia;

Gretche

Moffett Miss Lil Miss M

Durir Dutton'

The Festival

Seven

As S

honored by election to the office of Honorary President for one year. He was further honored by election to Honorary Membership for life in the Society, as were other veteran members, Hiram Burkholder, Clyde and Dan Alban, Jackson. Mr. Thornburg has been associated with Society activities

during the past 65 years.

(Continued on page 35)

LESSONS IN ORCHARD CHEMISTRY

By E. D. WITMAN, Research Associate Ohio State University Research Foundation

ELEMENTAL SULFURS

Sulfur occurs in its elemental state in huge underground deposits especially around the Gulf Coast. It is used directly as it is recovered from these deposits for the manufacture of dusting sulfur and wettable sulfurs. Lime sulfur is a *compound* of sulfur and calcium and will be discussed in a later lesson. Flotation sulfur is a form of *elemental* sulfur recovered from various industrial gases.

Sulfur is insoluble in and not wetted by water. It melts to a liquid when heated slightly above the boiling point of water. When ignited it burns with a bluish flame, yielding sulfur dioxide, a colorless, choking, corrosive gas. One of sulfur's closest chem-

Dusting sulfur is made either by subliming crude sulfur to get finely divided "flowers of sulfur", or by grinding crude sulfur to a very fine powder. Dusting sulfur usually contains a few per cent of a conditioning agent or "breaker", such as talc, gypsum, etc., to

tains a few per cent of a conditioning agent or "breaker", such as falc, gypsum, etc., to prevent balling and make a free flowing powder.

Wettable sulfur is essentially the same as dusting sulfur except that it contains a little synthetic wetting agent so that it will readily suspend in water.

Flotation sulfur frequently contains soluble substances as it comes from the recovery process and these solubles must be removed by washing with water before a product satisfactory for use on plants is obtained. Flotation sulfur is usually sold in paste form.

er elemental sulfur somewhat like flotation sulfur in performance is produced by emulsifying method sulfur in water in the presence of a dispersing agent and then removing most of the water to form a paste or perhaps all of the water to produce a

Suffur is melted with bentonite clay, allowed to re-solidify and is then ground to a cowder to produce various proprietary products. When added to water such products form fine dispersions of elemental sulfur.

Sulfur is generally used for its fungicidal action although it is known to be insecticidal to some species. Its action as a fungicide is not exactly understood, but it is relatively certain that some vapor evolved from sulfur is responsible.

Sulfur is regarded as generally compatible with most insecticides except the oils.

AMERICAN FRUIT GROWER

ccording to 25,000 and buted on a idual fruit n plans to vers, F. F. ith the unnay accept late.

ce of the ch served 1867 when

ued on as

iety in orration and

ed at their recognize

fruit in

100 years

eting next

the Ohio

Deshler-

nd, Ohio,

ciety was

as further

Member-

ere other

ler, Clyde hornburg

activities

round

manu-

r and

eated

flame,

chem-

owers

con-

ins a

overy

oduct

form.

then

ice a

ducts

sectiit is

oils.

OWER

MISS NANCY ANDERSON

QUEEN SHENANDOAH TO RULE BLOSSOM FESTIVAL


The famed Shenandoah Apple Blossom Festival being held at Winchester, Virginia, on May 2 and 3, will have as its lovely queen this year Miss Nancy Anderson, daughter of Secretary of Agriculture and Mrs. Clinton P. Anderson.

As Queen Shenandoah XIX, the first queen since 1941, Miss Anderson will rule over an estimated 100,000 loyal subjects who are expected to gather at Winchester for the two-day celebration.

Seventeen-year-old Miss Anderson, whose home is in Albuquerque, New Mexico, is a student at Mount Vernon Seminary, Washington, D. C. An active participant in dramatics and glee club work, Miss Anderson has also served as president and vice-president of her classes.

The 19th Shenandoah Blossom Festival has as its top committee worker Mrs. B. B. Dutton, who has taken care of the details connected with the Queen and her court since the sixth Festival in 1929. Mrs. Dutton's assignment has included contacting thirty schools and colleges who annually send young ladies to the Festival; writing and arranging the coronation scripts; entertaining the Court, costuming, transportation and housing.

During the thirteen years of Mrs.
Dutton's stewardship the following queens were selected: Miss Mary Jo Matthews; Miss Suzanne Polard; Miss Priscilla Dacre Morton of Winchester, England; Miss Ames Washington; Miss Francoise May of Belgium; Mrs. John Hay Whitney; Miss Nella Veverka of Czechoslovakia; Miss Cornelia Larus; Miss Gretchen Thomson; Miss Adelaide Moffett; Miss Genevieve Garner; Miss Lillian Somoza of Nicaragua and Miss Mollie Foulks Lee,



PESTMASTER DDT Insecticide and Fungicide Formulations give closer control over destructive insects that attack your apples, peaches, pears and grapes.

SCIENTIFICALLY dORRECT PREPARATIONS. PESTMASTER 50% DDT Wetable Powder, PESTMASTER Fruit and Garden Dust, and PESTMASTER Fruit and Garden Spray, have been thoroughly field tested and approved by various agricultural colleges, experimental stations and the research department of the Michigan Chemical Corporation.

KILLS. Codling moths, green aphids, Japanese beetles, oriental fruit moths, peach tree borers, rose chafers, berry moths and certain other insects that come in contact with fruit or leaves are controlled with scientifically correct PISTMASTER DDT Formulations.

EASY TO USE. PESTMASTER 50% DDT Wettable Powders are compatible with wettable sulphurs, Bordeaux mixtures,

fixed or proprietary coppers, summer oils, nicotine sulphate, fixed nicotine and DN compounds. Write us today for complete instructions regarding the use of PESTMASTER products for insect control on fruit, vegetables, and livestock. Limited number of dealerships still available.

MICHIGAN CHEMICAL CORPORATION

SAINT LOUIS, MICHIGAN



You don't have to be BIG

to have a profitable spray program

Ask your County Agent who, in your vicinity, offers spraying service on a co-operative or laborexchange basis . . . someone hav-ing the equipment, manpower and know-how to give you a biggrower program of insect control.

Alorco Cryolite will figure frequently in that program, because growers have proved it's an economical and dependable control for many chewing insects. It contains 90% chemically refined sodium fluoaluminate, the effective killing

Get all the details on Alorco Cryolite from your local dealer, or write

ALUMINUM ORE COMPANY

Subsidiary of Aluminum Company of America 1737 Gulf Bidg., Piersburgh 19, Pa.



LOOKING FORWARD TO THE 1947 CROP

(Continued from page 11)

considerable development during summer and late fall and for several months or weeks prior to the time it

opens in spring.

We can be reasonably sure that differentiation into either a leaf bud or a flower bud occurs just before or very soon after the cells surrounding the active growing point prepare to enter the rest period. Different buds on the same tree cease growth activity and begin their rest at various times during the season. Those on spurs, or short shoots, may begin to rest before the end of May; others, on more vigorous growths or on terminal shoots, may remain in an active condition for several months. In most varieties of apples, the flower buds are formed at the ends rather than the sides of shoots. In mature trees a large proportion of terminal buds are at the ends of spurs. Since the spurs grow for only a few weeks before the rest begins, it is safe to assume that the majority of flowers are initiated early in summer.

Flower Forming Substances

This suggests that the factors which determine blossom bud formation for the following year's crop must exert their influence rather early during the present growing season. Appropriate internal chemical conditions of the tree tissues, especially those in the vicinity of the growing points, must be present even before we can see with the aid of a microscope that embryonic parts of flowers have appeared. These internal chemical conditions are referred to as flower-forming-substances and they include adequate reserves of carbohydrates, a minimum of nitrogen and other nutrient materials and also traces of hormones.

The prospects for a crop of apples in 1947 depend first of all upon a group of factors that favor the initiation, or that do not inhibit the development of flower parts, during 1946. It is not too early, therefore, to consider what, if anything, the grower can do during the next few weeks or months to help in getting the 1947 crop on its way. It must be recognized at the outset that weather and many other factors are largely beyond his control, but the little definite knowledge we have about flower bud formation in fruit trees may nevertheless be used to advantage.

At the present time, many apple orchards that were reasonably well cared for during 1945, but which had practically no fruits during that year, have many flower buds. It frequently

happens that an "off" year is followed by a heavy crop. An unusually heavy set this spring will definitely handicap flower bud production for 1947. It may also bring about a biennial habit of bearing in individual trees or orchards that could last for a number of years. This would mean that the even years, 1946, 1948, etc., would be "on" years and the odd years, 1947, 1949, etc., would be "off" years. Beyears. Beginning in 1912 and lasting for 22 years, the apple crop in New York State in the "on" years averaged about 35,870,000 bu., and for the "off" years, 19,293,000 bu. Since 1932, the fluctuations in annual yields have been less violent although we still have alternation between large and small crops.

Good Cultural Operations

The objectives of good cultural operations in the orchard in 1946 for those who are interested in regular bearing should be (a) to avoid too heavy a set of fruit, if this is possible and (b) to aim for a moderately good crop of high quality apples rather than a maximum tonnage. The following suggestions might be considered in this

1. Special efforts should be made before the buds open to do a thorough job of pruning. This will involve the elimination of undesirable bearing surface that might set fruit but which would not be so located as to produce reasonably large, high quality, well colored apples. Wood which is shaded due to crowded growth on the under side of the branches, or in the interior of the tree, obviously comes in this class. In some cases, vigorous, well exposed bearing wood found on the top of tall trees, might well be removed, since fruit borne there often involves too much trouble and expense for spraying and harvesting. This is especially true if the tall, umbrella-like growth interferes with the penetration of sunlight to vigorous lower branches more easily reached in spraying and harvesting.

During the past years, many growers have found it difficult to do all the pruning that they had planned. This year, with many trees heavily budded, is an especially appropriate time to do more than the normal amount of pruning. Emphasis should be placed on the removal of the weak wood, leaving as much as possible of the vigorous bearing surface that is well exposed to

In mature trees, it is more important to maintain the existing bearing

(Continued on page 24) AMERICAN FRUIT GROWER

grapefi the re method in the c

Earl are in . first of the tre 'picker the tree nosed state in only p tangeri with a bruise Juice spector

grove a

up to r

These

the sta

by the to assu for the mate co in grov neries e and siz certifica for all leave th Dr. P (picture argest

in the s vidual gerine that of which i packed operativ the citr tangerin block co

Dr. P structiv the frui to impre in the i greater marketi

B One native t in the o the Der

Ohio A tion at growing neighbo

market The h we still

large and

ny growdo all the led. This y budded, ime to do c of prunlaced on d, leaving vigorous xposed to

enetration

branches

ying and

e imporg bearing

ROWER

NATIONWIDE FRUITS

(Continued from page 13)

grapefruit. Today's tangerines are the result of completely modern methods of citrus growing unknown in the days of Columbus.

Harvest Time

Early in the fall, trained pickers are in Florida groves to remove the first of the ripened tangerines from the trees. Belying their name, the "pickers" do not pick the fruit from the trees, but snip it off with bluntnosed clippers. Since growers and state inspectors cooperate in sending only perfect fruit to market, each tangerine must be expertly clipped with a short stem in order not to bruise others during shipment.

Juice tests are taken by state inspectors right under the tree in the grove and only fruit that measures up to maturity standards is picked. These standards are established by the state legislature and supervised by the Florida Citrus Commission, to assure the highest standards both for the citrus industry and the ultimate consumer. Some 250 inspectors in groves, packing houses and canneries enforce the maturity, grading and sizing standards and issue the certificates which must be secured for all citrus fruit before it may leave the state for northern markets.

Dr. P. Phillips of Orlando, Florida (picture on page 13), is one of the largest individual growers of citrus in the state, and is the largest individual tangerine grower. His tangerine tonnage runs second only to that of the Florida Citrus Exchange which is a cooperative handling the packed fruit from nearly fifty cooperative houses scattered through the citrus belt. Dr. Phillips has one tangerine grove alone in a single block covering a square mile of land.


Dr. Phillips is a progressive, constructive operator who believes in the fruit he grows, and does his best to improve handling conditions within the industry so as to permit still greater increases in production and marketing capacity for tangerines.

NUIS

Black Walnuts Available

One of the most valuable trees native to Ohio is the black walnut, in the opinion of Robert R. Paton of the Department of Forestry at the Ohio Agricultural Experiment Station at Wooster, Ohio. It is found growing in many parts of this and neighboring states and finds a ready market for many uses.

The typical habit of the walnut (Continued on page 27)







5 Pick-ups For The Farm

6½' pick-up body on ½-ton . . . 7½' pick-up body on 34-ton . . . 7½' pick-up body on 1-ton . . . 9' pick-up body on 1- and 1½-ton. All bodies are 48½" wide; sides 17" high to top of flare.



Stakes and Platforms, Too!

7½' and 9' stake bodies on 1-ton . . . 9' and 12' stake bodies on 1½-ton . . . 9' and 12' stake bodies on 2-ton. Chassis, with cab, available to take 14' bodies; cab-over-engine models also available where shorter wheelbases are desired.

You don't farm very long without realizing that one of the best ways to increase farm income is to reduce farm costs.

That's why farmers rate economy of operation right along with dependability when asked why they stick to Dodge Job-Rated trucks year after year. And that's why this year—as new trucks again become available—more and more farmers are switching to trucks built to fit farm jobs, and thus save money on the job!

If you're not too busy to talk about saving money—ask your Dodge dealer why a truck built to fit the job will do a better job on your farm, save you more money, and last longer!



LISTEN TO THE MUSIC OF ANDRE KOSTELANETZ, WITH FAMOUS GUEST STARS-THURSDAYS, CBS, 9 P.M. E.T.

THE 1947 APPLE CROP

(Continued from page 22)

CTOSS

But

velop

the to

bees help

well

of go

two

bloom

pecia only

the c

heav

linati

numl

velop

of co

Inne

these

draw

the to

a red

which

bohy

neces

serve

hand

are a

to re

thinn

late

grow

in br

after

thinn

sider

ment

indic

spone

ment

recon

charc

never

with

to se

Mate

ful to

shoul

that

and I

Man

recal

the f

latter

neigh

own.

defol

seaso bloom

Th

surface in good vigor than it is to try to correct any faults due to previous years' pruning. In the younger trees, on the other hand, the grower should keep in mind that the full bearing surface has not yet been attained, and more attention might well be given to the removal of crowding and interfering growth with the object of developing a strong framework of well placed limbs along a main leader, with properly distributed secondary branches.

Thorough Spraying

2. In most orchards throughout the eastern part of the country, the chances for scab in 1946 are very great because of the carry-over of heavily infested leaves from the bad scab year of 1945. Special attention should therefore have been given to the proper timing and thorough spraying with appropriate materials during the delayed dormant period to prevent primary infection. Pruning done in such a way as to facilitate the penetration of spray to all parts of the tree will help in this respect.

Fertilizer Application

3. The fertilizer program in most well cared for orchards should not be stepped up. As many growers have learned from experience, an abundant supply of nitrogen, applied before bloom tends to increase the set of fruit. Pruning likewise favors a heavy set. Where the trees have been in good vigor for the past few years and where the fertilizer program has been on the liberal side, a slight reduction in the rate of application might be desirable especially if heavy pruning has been done. Perhaps 1/3 or 1/2 of the normal amount would suffice where the 1945 twig growth was very good. This reduced rate is recommended also in those cases where the red color on the apples in previous years was short of the requirements of Fancy or No. 1 grade.

A delay in time of application of nitrogenous fertilizer until after bloom might be desirable this year. The object of such delay would be to obtain the invigorating effects of fertilizer on growth without over-stimulation of the set of fruit. There is, of course, the danger of interfering with the development of red color if too much nitrogen is available during the latter part of the growing season. Where early cultivation is practiced less nitrogen is usually required than in the case of sod orchards, but the cover crop should be established early.

AMERICAN FRUIT GROWER

CROP

n it is to try to previous unger trees, ower should bearing surtained, and ill be given

ll be given g and interbject of deork of well ain leader, secondary

oughout the ountry, the e very great of heavily bad scab tion should en to the th spraying during the to prevent ing done in

he penetra-

of the tree

ion

m in most ould not be wers have n abundant ied before the set of ors a heavy en in good and where een on the tion in the e desirable has been the normal e the 1945 . This reed also in olor on the

ication of attil after this year. ould be to cts of ferver-stimu-There is, interfering d color if ble during a season. practiced aired than s, but the hed early.

ROWER

s short of

or No. 1

Cross Pollination

4. Many varieties of apples require cross pollination for a satisfactory set. But too much cross pollination may cause too many fruits to begin to deyelop. This early set tends to throw the trees into alternate bearing. Where bees are brought into the orchard to help in transferring pollen, it may be well to remove the hives after one day of good pollination weather, provided two or more varieties have been in bloom during that time. With weather and other conditions favorable, especially when many bees are present, only a few hours are required to do all the cross pollination necessary for a heavy set of fruit. Any additional pollination would simply increase the number of flowers that begin to develop into fruits, many of which will, of course, be eliminated during the June drop. In the meantime, however, these extra number of fruits will have drawn heavily on the food reserves of the tree. The heavy set also results in a reduction of the early leaf surface, which limits the manufacture of carbohydrates, and therefore prevents the necessary accumulation of food reserves which must be present for the initiation of the 1947 blossoms.

Hand Thinning


5. Where the set is very heavy, hand-thinning as soon as the fruits are a few weeks old is almost certain to result in larger apples. Such handthinning, however, usually comes too late in most orchards under eastern growing conditions to do much good in bringing about annual bearing year after year. The possibility of blossom thinning by spraying should be considered in this connection. Experiments under way on this problem indicate that individual varieties respond differently to the same treatment and that as yet no general recommendation applicable to all orchards can be made. Some growers, nevertheless, may want to experiment with at least a part of their plantings with the object of having some fruit to sell in 1947 as well as in 1946. Materials that are known to be harmful to bees when applied during bloom should be avoided.

Thus far we have considered trees that were reasonably well cared for and had fairly healthy foliage in 1945. Many fruit growers, however, may recall the rather sad appearance of the foliage on apple trees during the latter part of 1945 in some of the neighbors' orchards if not in their own. Too many orchards have been defoliated so early during the past season that they will have little or no bloom in 1946. The short growth on

(Continued on page 26)



Why Bus Lines specify Champion Spark Plugs



THEY'RE DEPENDABLE!

The bus lines of America are indispensable to our transportation system. Like farmers, they are experts on maintenance of equipment because efficiency and economy are absolute essentials to success in their business-no factor is too small to escape detailed study. That is why most bus lines with the best maintenance records specify dependable Champion Spark Plugs. This is another example of preference for Champions by experts—substantial evidence that they're better for all farm engines, in cars, trucks, tractors and stationary units.

Champion Spark Plug Company, Toledo 1, Ohio.



FOLLOW THE EXPERTS . . . DEMAND DEPENDABLE CHAMPIONS

THE 1947 APPLE CROP

(Continued from page 25)

such trees is likely to be very spindly and the buds quite small.

Where the foliage remained green with relatively little scab until about the first of September, there may be some flowers present even though the trees are weakened. When the buds open, there will probably be only three or four flowers in a cluster instead of five or six, and the spurs will have small leaves, because of low reserves of stored food. The flowers will also be handicapped in setting a crop because of the lack of carbohydrate materials. With especially favorable pollination conditions, some fruit may be formed but it is very unlikely that the weakened trees will require any thinning.

Handicapped Orchards

How should we manage these handicapped orchards so they can produce a good crop in 1947, if not this year? Apple trees often show a remarkable ability to recover from the effect of a poor growing season provided the buds and the wood are not damaged by cold during the winter. Trees that have little or no fruit in 1946 should naturally develop an abundance of bloom for the 1947 crop, especially if they are kept free from insects and diseases and are otherwise given reasonably good culture. The following practices are important:

Practices to Follow

1. Thorough pruning to eliminate all dead or weak wood is the first step in rehabilitating the neglected orchard. The same procedure should be followed as previously suggested for the more vigorous trees. More wood, however, will probably have to be removed from the weak trees. Flower buds that may be present will be found on the well exposed portions of the tree. Such wood should be saved, if at all possible.

2. Since most of the defoliation last year was caused by scab, special efforts should be made to prevent another infection this year. It is well known that scab is likely to get on the exposed leaf surface soon after the buds break, and even before the flowers are open. No chances should be taken in "cutting corners" in spraying the orchards which were badly infested in 1945, even though there is little fruit in 1946.

3. Not much more than the normal amount of fertilizer should be used where the trees have no crops to support. Too much nitrogen could easily result in heavy sucker growth. The pruning, too, will help invigorate the trees.

might cial at qualit.
The the 1sh higher apples bear might bearing crop is practic removes surface at the cial at the surface at the cial at the cial

aimin

These

numb

vidua

crop :

while

Wi

1946.

oppor

severa

their

probal

inclina

the be

is to woods singly is ran stands work specie

It is be pla be set are fa assure tinuou

timbe
Pla
seedlii
have
outdo
factor
econo
still a
Nurse
cents
proxii

Any planti should Woosi tion. metho other of cha

no see

MAY

E CROP

page 25)

e very spindly

emained green ab until about there may be en though the hen the buds be only three ster instead of urs will have low reserves wers will also ng a crop becarbohydrate ally favorable ome fruit may

chards

unlikely that

1 require any

nanage these so they can 1947, if not often show recover from owing season the wood are ring the winle or no fruit y develop an he 1947 crop, pt free from are otherwise culture. The portant:

low

to eliminate the first step cted orchard. ould be folggested for More wood, have to be rees. Flower will be found tions of the e saved, if at

defoliation scab, special prevent an-It is well y to get on soon after before the nces should s" in sprayre badly ingh there is

the normal ald be used ops to supcould easily owth. The rigorate the

GROWER

Concluding Remarks

With the prospects at the present time for a large crop of apples in 1946, the consumers will have greater opportunity than they have had for several years past to discriminate in their purchases of fruit. They will probably have enough money and the inclination to pay higher prices for the better grades. Growers, therefore, might find it a good policy to pay special attention to the production of high quality apples.

The cultural practices suggested for the 1946 season, should result in a higher percentage of better grades of apples. If the trees are allowed to bear too heavily this season, they might easily be thrown into a biennial bearing habit, which means a small crop in every other year. The most practical way of avoiding this is to remove undesirable excess bearing surface by judicious pruning, and by aiming for a moderate set of fruit. These practices will tend to reduce the number but increase the size of individual fruits that develop. A good crop may thus be produced in 1946 while flowers are being initiated for the 1947 crop.

NATIONWIDE FRUITS

(Continued from page 23)

is to grow scattered through the woods in mixture with other trees, or singly in fields and fence rows. It is rarely, if ever, found in pure stands. This characteristic tends to work to the disadvantage of the species, for cutting the trees may eliminate the possible seed crop in that vicinity.

It is important that more walnuts be planted in Ohio, and they should be set out wherever site conditions are favorable. Such a program will assure future generations of a continuous supply of this valuable timber.

Planting may be done either with seedlings or nuts, provided the latter have been stored in damp sawdust outdoors. Both methods are satisfactory and the latter is much more economical. Nuts for planting are still available at the State Forest Nursery, at \$2.00 per bushel, or 75 cents per peck. A peck contains approximately 250-300 nuts. There are no seedlings available this spring.

Any Ohio grower interested in the planting of walnuts for reforestation should write to the State Forester, Wooster, Ohio, for further information. Instructions as to planting method, selection of the site, and other information will be sent free of charge, upon request.



"U.S." SAVOY COAT. Double backed for extra service. Heavy lining, corduroy edged collar, 11 ventilating outlets.

"U.S." SQUAM HAT. Reinforced brim. "U. S." RUBBER ARCTICS. Scientificstrength compounds at points of stress-buckles firmly anchored. 4 or 5 buckle.

"U. S." RUBBER BOOTS. Leaktested under water by internal airpressure-"tire tread" strength in sturdy "CLEATRED" soles—rocker last for walking comfort.

"U. S." WORK RUBBERS.





You can count on this . . . "U. S." KNOWS HOW TO MAKE WATERPROOF CLOTHING. The science and craftsmanship that build "U. S. Clothing and the quality of rubber compounds used are backed by 100 years of "U. S." experience.

Sold only through Industrial and Retail Stores

"U. S." ROYAL RUBBER FOOTWEAR



"U. S." INDUSTRIAL RAYNSTERS

Serving Through Science -

UNITED STATES RUBBER COMPANY

1230 AVENUE OF THE AMERICAS . ROCKEFELLER CENTER . NEW YORK 20, N. Y.



BUY THE BEST IN STEELS!

Planet Jr.

Farmers have learned that it's smart to look for the name Planet Jr. when buying Steels! Check these points and you'll see why "Planetized" Steel Shapes are famous for their amazing wearing qualities.

- Steel rolled especially to Planet Jr. specifications.
- Painstaking attention to smallest details of design to assure maximum efficiency.
- Special hardening operations

 under scientific control—for
 best wearing qualities.
- Proper tempering for still greater toughness-without brittleness.
- Special finish for better scouring.

All this adds up to a long, useful life for Planet Jr. "Planetized" Steel Shapes!

S. L. ALLEN & CO., Inc. 3439 N. 5th Street, Philadelphia 40, Pa. Makers of Planet Jr. Tractors Catalog on Request

Planet Jr.



WRITE for CIRCULAR

Wettable powder in strengths of 25% or 50% DDT. Use as water suspension spray. Kills codling moth, apple leaf hopper, Oriental fruit moth and certain other insects on fruits.

CHIPMAN CHEMICAL

Dept. K, Bound Brook, N.J.

ORCHARD POLLINATION

(Continued from page 10)

mendation under such conditions. Such topworked trees should bloom in 4 to 5 years if the grafting is in branches not more than 1 to 2 inches in diameter.

In the meantime, while waiting for such grafted trees to begin to bloom, he can use hand pollination, as discussed in another article in this issue of AMERICAN FRUIT GROWER, or he can bring in branches of pollinating varieties.

Pollinating "Bouquets"

The latter method has been proved effective if well done, and where bee activity is ample. It must be emphasized that many pollinating "bouquets" must be used, and they should be of large size. Generally pruning of the pollinizer varieties should be postponed to the bloom stage. Then a large crew must be put to work to prune the pollinizers and haul the branches to the orchard where they are needed. They must be kept from drying out-either by keeping them in barrels of water while being hauled, or by covering them with wet canvas and getting them placed in the orchard block within an hour or two after cutting. The larger the branches cut, the longer they will remain fresh and suitable for pollination purposes.


Suspend Above Ground

Fairly large "bouquets" should be used—as many branches as can be put in a 3-gallon pail at each place. The best location is suspended above ground between the trees. If overhead spray pipes are installed some of the pails may be suspended from them. Tripods made by lashing or bolting props together, with the pail suspended below the point of fastening, may be used to keep the pollinating bouquets up in the air. It is the general experience of growers that bees will visit these pollinizers better if they are even a few feet off the ground, rather than set on the

Branches 3/4" or more in diameter should generally be used. Place them in the pails so the cut end is under water—and keep water in the pails so the blossoms will last several days.

As to the number of bouquets to bring in, it seems that "the more the better" applies. A minimum for good effectiveness would be a large bouquet for each 4 trees, putting the bouquet in the square between them. The bringing in of bouquets should start as soon as bloom is beginning

(Continued on page 29)



ORG

creas

are of

Ev

the of

the P

In the

this i

and n

move

mean

The chandle

domes

them

ing st

spray

the be

The

dling

are m

ton, b

partm

move

or thr

al sta

bees t

new lo

are op

impor

coloni

size. A

ly one

more e

coloni

tremer

is par

standp

The o

premii

Bees f

will r

weak

"Ore

served more

than o

work

richest

change

blosso

a rich

sugar.

ple, co

conten

pete di

tation

is abu

around

bloom,

MAY,

"En

"In

COPPER SULPHATE

SAVES LABOR ... Mixes easily in spray tank. Eliminates stock solution, minimizes handling!

SAVES TIME . . . Dissolves rapidly. No waiting for it to go into solution!

SAVES MONEY... Dissolves thoroughly—no waste or sediment. Gives accurate control and greater safety in your mixtures—thus giving better protection and increased yields.

• FREE Send postcard today for val.
uable free booklet —"Bordeaux Mixture
— Its Efficient Preparation and Use."

YOUR DEALER can also supply Triangle Brand Copper Sulphate in: LARGE CRYSTALS, SMALL CRYSTALS, GRANULATED and SUPERFINE for regular Bordeaux Mixtures; also Monohydrated for Copper-Lime dusts. The oldest and the best known brand! Standard for over 50 years.



AGICIDE SPRAY BASE

For control many fruit insects NO HARMFUL RESIDUES

Agicide Gold Dot Spray Base

(50% D.D.T.)
For control codling moth
Write for name nearest dealer

AGICIDE LABORATORIES, Inc.

AMERICAN FRUIT GROWER

RDEAUX
S
GLE

PHATE RE!

g! issolves rapidinto solution!

es stock solu.

Dissolves or sediment and greater—thus giving reased yields.

today for val. eaux Mixture on and Use."

o supply Triulphate in: ALL CRYSand SUPERux Mixtures; Copper-Lime best known 50 years.

PR SAW

Fastert

Fast

insects
UES
ray Base

dealer IES, Int. Wiscensin

GROWER

ORCHARD POLLINATION

(Continued from page 28)

to open, with the number being increased up to the time most blossoms

Even with pollinating varieties in the orchard it is still necessary to get the pollen carried to the blossoms. In the case of these self-sterile fruits this is entirely the work of insects, and mainly of free flying insects that move rapidly from tree to tree. This means mainly bees of various types. The only one of these types we can handle or in any way control is the domestic honey bee. We can move them into the orchard when blooming starts, and remove them before spraying with materials poisonous to the bees begins.

Bees for Pollination

The following comments on handling bees for orchard pollination are made by Mr. James I. Hambleton, bee specialist of the U. S. Department of Agriculture:

"In general, it is preferable to move the bees into the orchard two or three days in advance of the actual start of bloom. This gives the bees time to become settled in their new location by the time the flowers are open.


"Emphasis should be placed on the importance of obtaining strong colonies. Colonies vary greatly in size. A strong colony, and particularly one rearing brood, may be much more effective than two or three weak colonies. Bees rearing brood have a tremendous appetite for pollen. This is particularly important from the standpoint of orchard pollination. The orchardist should insist on a standard of strength in renting colonies and should be ready to pay a premium for the strong colonies. Bees from strong colonies will work under more adverse conditions and will range wider than those from weak colonies.

Competitive Plants

"Orchardists have sometimes observed that the bees seemed to work more on other blossoming plants than on fruit blossoms. Bees tend to work on the blossoms that are the richest source of nectar and will change quickly from one type of blossom to another if the latter bears a richer nectar-that is higher in sugar. Mustard blossoms, for example, contain about the same sugar content as apple blossoms, and compete directly with them for the visitation of pollinating insects. If there is abundant mustard bloom in and around the orchard during fruit bloom, the situation calls for more

(Continued on page 30)





the No. 5 the most popular tractor mower ever built.

Working with any make of tractor, the No. 5 Power Mower cuts up to 35 acres per day. Simple, durable, easy to attach and detach, it's a tractor mower through and through. Strong, well-braced frame . . . enclosed drive which runs in oil . . . safety spring release hitch and slip clutch on power shaft to protect entire mower against breakage . . . high, easy lift . . . caster wheel and hinged drawbar carriage to insure maximum flexibility for all ground conditions . . . accurately fitted, long-lived cutting parts . . . simple, easy adjustments to keep the No. 5 working like new down through the years—these and many other quality features combine to give you the best tractor mower money can buy.

The No. 5 Power Mower is built with 7-, 6-, 5-, and $4\frac{1}{2}$ -foot cutter bars. See your John Deere dealer for further information.



JOHN DEERE, Moline, Illinois



ou can greatly increase the kill of orchard pests with Pratt's 25% DDT—soluble DDT in its most effective form. It leaves a microcrystalline deposit on the foliage which increases toxicity. 2 pints in 100 gallons of water is especially effective in controlling codling moth.

Kills potato beetle, thrips, leaf roller, flea beetle, Japanese beetle, oriental peach moth and many caterpillars. Pratt's 25% D D T makes a powerful and long-lasting residual fly spray for spraying barns, dairies, kennels and other farm buildings. Order today from your dealer.

B. G. PRATT CO., 163 RIVER ST., HACKENSACK, N. J.

Mfr's of Scalecide, Para-Scalecide, 25% DDT (liquid), 50% DDT (dry-wettable), D-X Spray, Surfispray

Protect Your Profits...



Get complete orchard protection. Power agitator and spiral feeding mechanism prevent bridging and deliver a full, even flow of dust to nozzle regardless of type dust used. Can be operated by one man. Built of welded steel plate for maximum strength with minimum weight. Economical, trouble-free performance.

HOWRY-BERG STEEL & IRON WORKS
2949 N. Speer Blvd., Denver 11, Colo.

NORTHERN NEWLAND GROWN PLANTS Wholesale Prices

		100	250	500	1000
Streamliner New	Supreme Everbearer	\$6.00	\$14.00	\$25.00	\$40.00
Minnesota 1166	Everbearer	3.00	6.50	12.00	18.00
Gem Everbearer	Crimson Beauty	2.00	4.50	8.00	16.00

RASPBERRIES 1 YR. PLANTS

Sunrise or Latham	Red Rasp.	8.00	18.00	35.00	70.00
Cumberland-Black	***************************************	4.00	9.50	18.00	32.00

Ask for CATALOG or ORDER DIRECT from Ad.

WESTHAUSER NURSERIES

(Box G

SAWYER, MICHIGAN

GROWING CONTRACTS

Gruit Trees

We can make growing contracts now for 1-Yr. apples, June Budded peach, or plum for Fall 1946 or Spring 1947 deliveries. Pears, cherries, prunes, 1-year peach, Fall 1947 and Spring 1948 delivery. Customers may, if desired, furnish buds or scions under a key number without disclosing to us or any one the name of variety. 2,000 smallest contract, maximum 100,000. Catalog free. Tennessee Nursery Co., Box 4, Cleveland, Tenn.



Write for Free Circular 12 KER-Ó-KIL MFG. CO. Gettysburg, Pa. or

Gettysburg, Pa. or Redwood City, Calif.

Have plenty of fruit to eat and sell this year. New MINN 1166 everbearing strawberry. Fruitin 60 davs till frost this Fall. Special Price 25-\$1.00, 50-\$1.50, 100-\$2.50, 1000 for \$20.00. Latham Red Raspberry 25 for \$2.00, 50 for \$3.00. New Streamliner everbearing strawberry 100 for \$7.00. Everything postpaid.

SOUTH MICHIGAN NURSERY
DEPT. FG2 NEW BUFFALO, MICH.

FARMERS WHOLESALE NURSERY

PAUL PATTERSON, Owner

A dependable source for true to name fruit trees. We specialize in commercial accounts, and exercise every care in keeping our varieties straight.

Write for prices.

P.O. Box 65

Smithville, Tennessee

You Get What You Buy

ORCHARD POLLINATION

(Continued from page 29)

THE

sect w

insect

tion a

eons

been'

plant

The

most

of po

lows i

ing it

the u

the p

plant'

es fro

and w

with the in

jacket

going

posits

plants

H

The

ambas

the A

Amer

ward

Canad

their

to fee

ing c

count

In

which

of ani

plants

of pol

tozoa

plant.

-the

meast

natur

are in

extre

that

every

deed

canny

form

paths

specif

agent

evide

famil

down

suma

throu

MAT

Me

bees or the mowing of the mustard if there is a question of adequate pollination.

Distribution of Stands

"The orchardist is often uncertain as to the best distribution of stands of bees. They are sometimes distributed singly throughout the orchard; at other times are congregated in rather large groups of colonies. Best results seem to be obtained when the colonies are distributed throughout the orchard not singly but in rather small groups of four or five colonies at a place. This reduces the work of looking after the bees, of distributing them and removing them from the orchard. interferes less with other orchard operations, and appears to be even better from the pollination stand-

"The minimum temperature for bee activity is about 50° F. Cloudy weather and winds also depress bee activity and limit the range of flight. Under such unfavorable conditions the importance of strong colonies is particularly apparent.

One Colony for Two Acres

"Under most conditions one strong colony for each two acres of orchard should be sufficient to insure adquate pollination provided weather conditions are at all favorable. During unfavorable weather, the more colonies used, the better."

This discussion has dealt mainly with apples. The problem with pears is similar. With sweet cherries and Japanese plums it is somewhat further complicated by the fact that a number of varieties are not only self-sterile but inter-sterile. The growers of these fruits must know the satisfactory pollinating varieties and provide for them in the orchard.

Transfer of Pollen

It is well also to keep in mind that even in self-fertile varieties pollen must be transferred from the stamens to the pistils. In good weather, but more particularly during bad weather, pollinating insects help in this essential step of transferring pollen.

If the fruit grower provides suitable pollinizers and has ample be supply, and if the weather is moderately favorable, a good crop of fruits should be secured. At least the grower will have done what he can to insure it.

AMERICAN FRUIT GROWER

LINATION

page 29)

the mustard adequate pol-

Stands

ten uncertain ion of stands metimes dishout the orare congree groups of seem to be onies are dise orchard not all groups of a place. This ooking after ng them and the orchard. ther orchard s to be even ation stand-

perature for F. Cloudy depress bee nge of flight. le conditions g colonies is

o Acres

as one strong es of orchard insure adeded weather orable. Durer, the more

dealt mainly n with pears cherries and mewhat furfact that a re not only terile. The must know ing varieties the orchard.

in mind that

llen

ieties pollen om the staood weather, during bad ects help in transferring

ovides suitample bet er is moderop of fruits st the growhe can to

GROWER

THE GREAT MYSTERY OF POLLINATION

(Continued from page 9)

sect would quickly starve; without the insect, the plant would lack fertilization and the species would perish. For eons this cycle of cooperation has been going on, unsuspected by either plant or insect.

Honeybee Most Active

The honeybee is unquestionably the most active gatherer and distributor of pollen. In its quest for nectar, the busy little creature virtually wallows in pollen dust. The hairs covering its body and appendages are of the utmost importance in picking up the pollen grains which cling to the plant's stamens. When the bee emerges from a flower, its head, neck, legs and whole body are so heavily dusted with the powdery yellow grains that the insect more resembles a yellowjacket than the familiar black bee. In going from flower to flower, it deposits this pollen on the pistils, thus accidentally pollenizing thousands of plants in the course of a lifetime.

Hummingbirds Carry Pollen

The humming birds are international ambassadors of pollination. Most of the American species winter in Central America and each year migrate northward to Mexico, the United States, Canada, Alaska and Labrador. In their long flight northward they stop to feed on plant nectar, and in so doing carry pollen from country to country along the way.

In lieu of the act of copulation which accounts for the perpetuation of animal species, nature has provided plants with the regenerative process of pollination. What the male spermatozoa is to mankind, pollen is to a

Methods of pollination are countless and while the end is the same -the fertilization of plants-the measures and subterfuges devised by nature to attract the pollenizing agents are ingenious and diversified in the extreme. To the layman it may seem that many of these methods leave everything to chance. This would indeed be the case were nature not so canny in placing temptation—in the form of nectar and pollen-in the paths of properly interested parties, specifically, the correct pollenizing agents. That the plan is foolproof is evidenced by the fact that all of our familiar flowering plants have come down through the ages intact. Presumably the ritual will continue through eternity.





TRUCK CROPS respond with speed and profit to 'AERO' Cyanamid ... agriculture's most useful form of nitrogen. This modern nitrogen fertilizer fits effectively into your crop production program because it:

- Provides 20.6% non-leaching mitrogen.
- Provides 70% lime to maintain he calcium in your soil and avoid
- Gives a long lasting residual effect proved to be the most beneficial to soil of any nitragen fertilizer
- 1 Improves tilth and friability of the land through development of humus when plowed down with cover crops or crop refuse.

Write for special recommendations covering your truck or fruit crops. American Cyanamid Company, Fertilizer Division, 30 Rockefeller Plaza, New York 20.

yanamu

AGRICULTURE'S MOST USEFUL FORM OF NITROGEN

FREE BOOKLET



Write today for the new booklet describing Es-Min-El, the soil mineralizer that makes it possible for you to grow quality fruits and vegetables! Put minerals and vitamins into the food you grow by using Es-Min-El.

TENNESSEE CORPORATION

ATLANTA, GEORGIA. LOCKLAND, OHIO

HAMILTON "Spray Guns that pay"

Your Spray Program is no better than your Spray Gun

See the New Style Hamilton Guns with Controlled Streamline

W. L. HAMILTON & CO.

Seven Models A GUN FOR EYERY PURPOSE Send for FREE Catalog

BROWER Weed Burner and Disinfecting Torch

Eansest, best way to kill weess—destroys seed and all. Ideal for disinfecting poultry and ho houses, barns, equipment. FAST—covers 1,000 sq. ft. per hour. Burns kerosene or distillate—4 gal. capacity (2 hour supply). Money-back Guarantee. Order from this ad.

DROWER MFG. CO. Day 7930 Quincy, III.

V DACVETC

By the thousand or by the carload THE MARIETTA FRUIT PACKAGE & LUMBER CO. MARIETTA, OHIO

Low Price OTTAWA "Buzz" Master

CLEARS LAND FAST! Powerful 6-HP motor with friction clutch for safe operation. Cuts down timber, brush and hedge; turn blade vertical and saw logs to length. Can be equipped to fell largest trees. Has clutch pulley for belt work. Fully guaranteed.

OTTAWA MFG. CO., 532 Brush Ave., Ottawa, Kans

Advertisement



From where I sit ... by Joe Marsh

Bird's-Eye View of America

A fellow took an aerial photograph of our town, and it makes the place look like Utopia.

Folks argued that the new firehouse would never look well beside the old Town Hall. But they harmonize perfectly from the air. One side of the railroad tracks looks as good as the other. All the different landmarks blend in nicely with surroundings.

From where I sit, there's a lesson in that photograph. A community's made up of different elements—people as well as landmarks. Some vote one way, some another; some enjoy a glass of beer and others don't. You might think there was a lot of reason for friction.

But it's all in your point of view. Get up high enough—see the community as a whole—and those little discords blend together into what we call America—a free, harmonious land. The differences only look big to people who see them from too close!

Joe Marsh

Copyright, 1946, United States Brewers Foundation

HAND POLLINATION

(Continued from page 14)

hest to

strong to polli

times C

tire tre

three p

linating

the oth

of the t

linate a

prising

seen th

son is 6

a day.

ence d

cious 1

to do

hand p

tain W

to Wis

BEEN

and if

fruit g

his tra

throug

whole a

ly coop

a bette

keeper.

truckir

arrang

of a r

beekee

200 co

long p

mercia

away.

strengt

bees s

could l

stung.

were r

were ir

helped

keeper

back as

through

the be

than s

to pay

all the

well s

service

out of

a good

tricts a

tain the

In one

coopera

200 co

by a sk

ment s

countie

package

MAY,

Alre

Sinc

ering pollen and selling it to growers. hire collectors to gather the pollen in commercial orchards. Each collector has a quart jar with a medium mesh screen fastened over the top, replacing the cover. Usually two to three flowers are removed from a flower cluster leaving the central flower (king blossom) and one or two lateral flowers. The anthers are taken when the flowers are in the plump or "balloon" stage, which is just before the petals have spread apart. At this stage the anthers are still closed. The flowers may be open, but it is important that the anthers have not dehisced. A flower is removed from the spur, rubbed over the mesh screen so that the anthers-fall in the jar. Blossoms are collected from all parts of the tree. In order to keep the pollen in good condition, it is measured frequently, and it is then kept in the shade until taken to the curing room. The gathered anthers are spread very thinly on large paper trays that are placed on racks or shelves. The drying room should be preferably kept at 70°F. During the drying process the anthers dehisc and release the pollen. After the pollen is dry, it is packed in ounce quantities in small containers and is then ready for shipment or use. Some dealers use a carrier with the pollen so that more blossoms can be pollinated with the same amount of pollen. Some of them mix the carrier with the pollen before sending it out; others send it and pollen separately and the grower mixes the pollen as he thinks necessary. As a result of careful investigation it is known that the pollen remains viable for two to three weeks at least from the time of curing. The pollen is usually tested for germination just before it is sent out. The purchaser of pollen should store it in a refrigerator or keep it at about 35°F. It is also very important to keep it dry.

Use Artist's Brush

When applying the pollen to the flowers, persons doing the pollinating attach the container with a small amount of pollen to their clothing so that it can be easily and conveniently reached with a brush. A long-handled artist's brush of which the diameter of the bristle portion is about 1/4 inch is first dipped into the pollen, excess, pollen is shaken off, and the brush applied to the blossom. The central blossom in the cluster is usually pollinated shortly after the petals have unfolded. It is important that pollen is placed on all five stigmas. The number of blossoms pollinated on a tree varies with the grower. It seems

AMERICAN FRUIT GROWER

TION

re 14)

to growers, ne pollen in ch collector dium mesh p, replacing three flowwer cluster (king blosral flowers. n the flow-"balloon"

the petals s stage the he flowers ortant that ed. A flowour, rubbed hat the anms are colne tree. In good conuently, and until taken athered anly on large on racks or should be During the dehisc and ne pollen is quantities then ready me dealers len so that nated with . Some of the pollen ers send it the grower inks necesl investigapollen reree weeks

portant to len to the pollinating n a small clothing so nveniently ng-handled e diameter out 1/4 inch len, excess, the brush

uring. The

r germina-

out. The

store it in

at about

he central sually poletals have that pollen mas. The ated on a . It seems

GROWER

best to pollinate every fourth or fifth strong spur. There is some tendency to pollinate too many blossoms. Sometimes one person takes care of an entire tree, and at other times, two to three people work on a tree, one pollinating the lower branches, while the others use ladders to get the tops of the trees. The time required to pollinate a medium-sized tree seems surprisingly short to those who have not seen the operation before. One person is expected to pollinate 15-20 trees

Since Wisconsin growers experience difficulties in getting the Delicious to set fruit, we are planning to do some experimental work on hand pollination this spring to ascertain whether this method is adapted to Wisconsin conditions.

BEEKEEPERS' VIEWPOINT


(Continued from page 16)

and if the ground is soft and wet, the fruit grower should lend a hand with his tractor to get the bees distributed through the orchard. In fact, the whole arrangement calls for neighborly cooperation. Under present labor conditions, the fruit grower may have a better supply of labor than the beekeeper. He might even help with the trucking. An outstandingly successful arrangement on a large scale was that of a northern Michigan commercial beekeeper who rented approximately 200 colonies of bees per year for a long period to pollinize a large commercial orchard nearly 100 miles away. The beekeeper guaranteed strength of colony and prepared the bees so that inexperienced helpers could handle the hives without being stung. He made certain that the bees were not going to starve while they were in the orchard. The fruit grower helped move the bees from the beekeeper's apiaries to the orchard and back again and helped distribute them through the orchards, and still paid the beekeeper a better rental price than some fruit growers are willing to pay when the beekeeper furnishes all the service. The fruit grower was well satisfied with the pollinating service and the beekeeper got enough out of it to make it worth while to do a good job.

Already fruit growers in some districts are finding it impossible to obtain the desired number of rental bees. In one county, the fruit growers are cooperatively buying approximately 200 colonies of bees to be operated by a skilled beekeeper. This arrangement should be satisfactory. In other counties, fruit growers are buying packages of bees to be operated by a

(Continued on page 35)





221 N. LaSalle Street, Chicago I, Illinois 636 California Street, San Francisco 8, California





PICKING BAGS

For Apples, Peaches, Pears Save your fruit Speed your picking
Write for folder JOHN BACON CO. Gasport, N. Y.

HERE'S YOUR BARGAIN

in CUSHION MUMS. For \$1.00 we send you at planting time our assortment of six colors hardy field grown plants. Colors bronze, rose, red, white, yellow, and red with yellow center. You will have hundreds of blossoms all fall. Mail your order today. order today.

RIVERSIDE NURSERIES, Dept. F1, Route 5, Elkhart, Ind.



OTTAWA Wood Saw FOR TRACTORS

Fast wood sawing, quickly pays for self. Easily moved while attached. Big blade. Free details. OTTAWA MFG. CQ. '832 Oak Ave., Ottawa, Kans

CANADIAN GROWN SEEDS - PLANTS - TREES REUSTRATED CATALOGUE FREE TOBE'S TREERY—NIAGARA-ON-THE-LAKE, CANADA



VATER SAVES TIME AND WORK

A BURKS Water System will give you running water where vou need it—Will help increase

milk and egg production—Put meat on hogs and beef faster—Cut chores and give you extra hours for field work.

BURKS WATER SYSTEMS

are famous for dependable service— longer life—and larger capacity. Only one moving part—self-priming—easy to install. Both deep or shallow well sys-

DECATUR PUMP CO., 10 Elk St., DECATUR 70, ILL.



OPPORTUNITY ADS

Only ISc a Word—CASH WITH ORDER. Count each initial and whole number as one word.

ADDRESS: AMERICAN FRUIT GROWER, 1370 Ontario Street, Cleveland 13, Ohio

BABY CHICKS

HELM'S PULLOBUM CONTROLLED CHICKS. HOLD-er Four World Records. ROP sired matings. Sexed Chicks. Free Brooding Bulletin. ILLINOIS HATCH-EBY, Metropolis, Illinois.

KEY, Metropolis, Illinois.

250-350 PEDIGREED SIRED BIG FLOPPY COMB
Type egg-bred "AAA" English White Leghorn Pullets
\$15,50. Unsexed \$8.50. Cockorels \$2.75. Four Weeks
0dd "AAA" Fullets \$25,50. 95% sex guaranteed. Catalog. MARTI LEGHORN FARM, Box A, Windsor, Missourt.

sourt.

SCHLICHTMAN'S U. S. APPROVED, PULLORUM tested chicks. Per 100 prepaid. Leghorns, Rocks, Rods, Orpingtons, Wyandottes, Minorcas, \$9.90. Assorted \$7.45. Pedigree sired and sexed chicks. Free catalog explaining 2-week replacement guarantee. SCHLICHTMAN HATCHERY, Appleton City, Missouri.

FDUCATIONAL

CORRESPONDENCE COURSES AND SELF-INSTRUC-tion books slightly used. Sold, Rented. Exchanged. All subjects. Satisfaction guaranteed. Cash paid for used courses. Complete information and 100-page illustrated bargain catalog FREE! Write—NELSON COMPANY, Dept. 2-65, Chicago 5, Illinois.

FERTILIZERS

AMMONIUM NITRATE—32.5% NITROGEN, JUNE DE-livery, 30 ton minimum cars. Suitable for side dressing corn, etc., top dressing pasture. Dealers wanted, SCHROCK FERTILIZER SERVICE, Congerville, III.

FOR SALE-EQUIPMENT & SUPPLIES

PORTABLE, GASOLINE, WESTINGHOUSE ELECTRIC Generating Units. 20 KW, 110 Volts, A.C. Slightly Used. \$995.00. ACE MACHINERY & EQUIPMENT COM-PANY, 115 W. Pratt Street, Baltimore I, Maryland. SPRAY GUN HOLDER. Enjoy Spraying with one hand. One arm sprayer. \$2.75 each delivered. APPLEWOOD SALES, Deerfield, Michigan.

Power Dusters for row crop and Orchard. New Flexible Spray Hose. Centrifugal Pumps and motors. Power and Gravity Conveyor. Bushel Turners. Bushel Cushions and Liners, Facing Plates and Shells, Irrigation pipes and systems compilete. Electric Shop welders, 250 Amp., and Air compressors. If it helps to grow or pack fruit TRY US. MICHIGAN ORCHARD SUPPLY COMPANY, South Haven, Michigan. Phone 252.

Haven, Michigan. Phone 252.

FOR SALE: CIDER PRESSES AND SUPPLIES. APple and peach Electro Magnetic Sizers. W. G. BUNKLES.

MACHINERY COMPANY, 185 Oakland Street, Trenton.

New Jersey.

New Jersey. DISINFECT WITH FIRE. WÖRLD'S FINEST AEROIL kerosene Torches destroy parasites, green weeds, seeds and all, instantly. 300,000 users. 4 gal. Tank, Burner and Hose \$22 express collect. 5 gal. \$37.25. Immediate shipment. Descriptive literature FREE. SINE EQUIPMENT, APG, Quakertown, Pa.

MALE HELP WANTED

OPPORTUNITY OF LIFETIME SUPPLYING DDT AND other profitable products to farmers. No experience or capital required. Must have auto and good references. Permanent. Write or wire MCNESS COMPANY, Dept. 705, Freeport, Illinois.

NURSERY STOCK

DEPENDABLE FRUIT AND NUT TREES, SMALL fruits, Ornaments, and General Nursery Stock Combined catalogue and Planting Guide free. CUMBERLAND VALLEY NURSERIES, INC., McMinnville, Tennessee. STRAWBERRY PLANTS — BLAKEMORE, MISSION-ary at 106 for \$2.00; 500 for \$5.00; Postpaid. 1000 for \$8.00. Erechearing at \$15.00 per 1000. Boysenherry thorn-less at 20c cach. JOHN LIGHTFOOT, Birchwood, Tenn.

less at 20c each. JOHN LIGHTFOOT, Birchwood, Tenn. IMPROVED BIJUEBERRIES. DELICIOUS U.S. GOV'T Hybrids large as grapes. 2 Yr. Plants 70 cents each, \$8.00 per dosen. 3 Yr. Bearing Age Plants \$1.35 each. \$15.50 dos. GEO. C. MORSE, Williamson, New York.

NEW STREAMLINER, EVERBEARING STRAWBERRY is the latest thing out. Berries enormous but never hollow. Bears 60 days after set to November. 3 crops in 18 months. (The Beauty Queen of Strawberries). Sweet, delicious, and firm. Prices: 25 for \$1.50; 50 for \$4.50; 100 for \$7.00; 250 for \$1.50. Postpaid. SOUTH MICHIGAN NURSERY, New Buffalo, Michigan.

for \$2.50; 100 for \$4.00; 250 for \$7.50; 1000 for \$11.20 Prepaid. SOUTH MICHIGAN NURSERY, New Modal Michigan.

GEM, AND MASTODON EVERBRARING THAT Prices: 25 for \$1.25; 50 for \$2.00; 100 for \$4.00; 100 for \$4.00; 20 for \$4.00; 100 for \$4.00; 20 for \$4.00;

CUMBERLAND BLACK RASPBERRY, LATHAM, Burred, Indian Summer, everbearing red, Sumise, but and red, Indian Summer, everbearing red, Sumise, but and red, Purple Raspberry, Thorniess Developer, Thorniess Developer, Thorniess Blackberry, Thorniess Blackberry, and Alfred Coreless Blackberry, Reduced States, 1988, 1989

Michigan.

Ss-DAY HYBRID SWEET CORN. THINK OF HAVING sweet corn to eat 58 days after planted. It can being done, with our seed. Prices: % 1b, 50e; Poul for 75e. Postpaid. SOUTH MICHIGAN NURSERY, IN DURAND, MICHIGAN.

Burialo, Michigan.

PEACH AND APPLE TREES LOW AS 20t. Plain
plums, cherries, nuts, berries, grapevines. Ide. Bragreens, shrubs, shade trees ow as 25c. Catalog Pra
TENNESSEE NURSERY COMPANY, Box 20t. Car-

ORCHARD FOR SALE

FOR SALE—YOUNG APPLE ORCHARD, ABOUT IN trees, commercial varieties, building-equipment, health, low price. FAIRVIEW ORCHARDS, W. REOG CLINGER, Prop., Milton, Pa.

ORCHARD SUPPLY BUSINESS FOR SALE

FOR SALE—REPAIR AND SALE OF FARM 11-chinery and Fruit Growers Supplies, last year's business over \$150,000. This combined with 55 acres young bears orchard showing a large return; nine room bearing home with all modern improvements; new two stay is x 60 sale and repair building with basement storas; lasm modern basement barn; four car garage; all buildings excellent condition; located heart of New York Fruit bat near Rochester; old established business with franciscincluded. Owner retring will sacrifice for \$75,000 win terms. GOODELL, Broker, Webster, New York.

PATENTS

NATIONAL TRADE MARK COMPANY, MUNIST Building, Washington, D. C.

PHOTO FINISHING

CENTURY DELIVERS WORK AND SERVICE THAT CAN't be beat. 8 exp. roll 25c with two professional alargements FREE. CENTURY PHOTO BERVICE 14 Crosse, Wisconsin.

Crosse, Wisconsin.

8 FINEST FOTOGRAVURE PRINTS FROM YOUR WILL
roll 25c, also free membership in Good Neighborn Plate
4 Tub which gets you 8x10 \$\frac{3}{2}\$ enlargement free. Strictly
honest deal. GRAPHIC CAMERA STUDIO, Hardwel,

PLANTS FOR SALE

PLANTS FOR SALE

SWEET POTATO PLANTS. NANCY HALL PORTO
RICO, Early Triumph and Southern Queen. Opengrose
and healthy, \$3.50 per 1000 postpaid. We manage
prompt shipment and absolute satisfaction. Ballit
RROS. PLANT FARM, Adairville, Kentucky.

SWEET POTATO PLANTS—NANCY HALL, PORTO
RICOS. Extra sweet, big, strong, vigorous, well resist.

200 \$1.00: 500 \$1.75: 1000 \$2.75; 5000 \$13.00. 1 C.

DELLINGER, Gleason, Tennessee.

SWEET POTATO PLANTS. NANCY HALLS—PORTO
RICARS. Satisfaction guaranteed. Strong, vigorous jains.

500 \$1.40: 1000 \$2.50; 3000 \$7.00. GUINS FLART
FARM, Sharon, Tennessee.

SWEET POTATO PLANTS. NANCY HALLS—PORTO
RICARS. 3000 bushels seed bedded, Strong, thrifty plant
guaranteed. Pay when received 100 50e; 300 \$1.50; 30

\$1.75: 1000 \$2.60: 5000 and over \$2.50 per 1000 RICS
PLANT FARM, Sharon, Tennessee.

RABBITS

The

GUARANTEED GIANT CHINCHILLA RABBITS. Pulific, Profitable. Most Valuable Fur. Markets Furnisat Details Free. WILLOW BROOK FARM, R2S, Scho-ville, Pa.

LIFETIME PAYING BUSINESS, LEARN COMMERCIAL Angors Wool Rabbit Raising, easily—quickly, Peticulars Free, ANGORA INSTITUTE, Box 6678, Sam. Oregon.

RAISE RABRITS, MAKE MONEY AT HOME. MAR-kets guaranteed. Booklet free. TAYLOR, Box 363-7.

RUBBER

RUBBER — YOUR RUBBER REQUIREMENTS ARE important—know where to get it when you want R Belts—rubber bands, boots, coats, gloves, boes, earl bulbs, aprons, budding strips. Get our Price Sheet and Catalog today, BROADWAY RUBBER MANUFACTURING COMPANY, Louisville 2, Kentucky.

SONGWRITERS

SONG POEMS WANTED. FOR MUSICAL SETTING. Send poem for immediate consideration. FIVE 57AS MUSIC MASTERS, 716 Beacon Building. Boston 8. Mag.

TREE BANDS

EARLY BIRD TREE BANDS CHEMICALLY TREATED.
Kills the Codling Moth. Send orders early. EDWIN E.
HOUSE, Saugatuck, Michigan.
BETA NAPHTHOL TREATED APPLE TREE BANDS
kill codling moth! Write for literature and prices. M.
A. KOELLER, Barry, Illinois.

RAISE TURKEYS THE NEW WAY. WRITE FOR FILE information explaining how to make up to \$3,000 in you own backyard. Address NATIONAL TURKEY INST-TUTE, Dept. 233, Columbus, Kansas.

FOR SALE—HYDRAULIC CIDER PRESS. 45 INCH Racks, Double Hydraulic Pumps \$450,00, FRUTH MARKET, 8. Perry, Napoleon, Oblo.

HAVE 950 SMUDGE POTS FOR ORCHARD USE FOR sale. For information, write LOUIS HAGA, Route 1, New Ers, Michigan. MICHIGAN NURSERY, New Burnalo, Michigan.
NEW MINN. 1166 EVERBEARING STRAWBERRY FOR folks who "Can't Eat Strawberries." Almost without acid. Sweetest of all. Large, dark red berries soon fill quart box. Profitable. Bears 60 days after set till November. 3 crops in 18 months. Prices: 25 for \$1.50; 50 FOR SALE—Immediate Delivery—New Apple Graders complete with polishers 400 and 800 bushels per day.

> "You sure picked yourself a job-tryin' to make that old car outwear those B. F. Goodrich tires."

The new B.F. Goodrich tires for passenger cars actually outwear prewar tires. The wider, flatter tread and special construction make these tires wear s-l-o-w-l-y and evenly. For your truck, tractor, and farm implements

there are special B.F.Goodrich tires backed by a policy of continued improvementpolicy that means more for every tire dollar you spend. Ask the B.F.Goodrich man to show you the new 1946 tires — and the new tubes that hold air longer.

An advertisement of B. F. Goodrich - First in Rubber

\$7.50; 1000 for \$7.50 URSERY, New Bursin, RBEARING STAY.

Always dependant
100 for \$3.00; 250 for
SOUTH MICHIGAN

ARING RED RAM.
in July and sens a
rerage family. Priss.
\$9.00; 100 for \$15.00.
NURSEBY, Now be-

RRY, LATHAM, REST ed, Sunrise, best only Dewberry, Thorses Blackberry, Rebest r \$5.00; 100 for \$1.00 RSERY, New Burlin,

THINK OF HAVING anted. It can and is 1/2 lb. 50e; Pound for AN NURSERY, New

OW AS 20c. PRANS, rapevines 10c. hers 25c. Catalog Fra.
NY, Box 202, Care

SALE HARD, ABOUT 1800 ling-equipment, HARDS, W. BROCK

ESS FOR SALE E OF FARM M. last year's business with room beauting in the room with room in the room

MPANY, MUNIST

NG

D SERVICE THAT

FROM YOUR FILM od Neighbors Photo ement free. Strictly STUDIO, Hartford, LE

Y HALL, PORTO Queen, Opengrown aid. We guarantee isfaction, BAILEY entucky.

Y HALL, PORTO
orous, well recei.
000 \$13.00. J. C.

HALLS—PORTO

HALLS-PORTO ong, thrifty plants 0c; 300 \$1.25; \$00 00 per 1000. RICE

RABBITS, Pac-farkets Furnished LM, R23, Selles-

EARN COMMER-ily—quickly, Par-Box 667B, Salen,

REMENTS ARE en you want it. oves, hose, gray Price Sheet and ER MANUFAC-ntucky.

ICAL SETTING. n. FIVE STAR Boston 8, Mass.

LLY TREATED.

TREE BANDS

TE FOR FREE \$3,000 in your RKEY INSTI-

GROWER

OPPORTUNITY ADS

(Continued from page 34)

USED AUTO PARTS

HAVING CAR TROUBLE? New, USED, GUARANised auto, truck parts save money. Transmission specialised secretic needs; immediate reply. VICTORY, 2936
is North Western, Chicago 18, Illinois.

WELDING

MAGIC ELECTRIC WELDER, 110 Volt AC-DC; welds, brass, solders, cuts all metals, easy to use; full directions of the complete with power unit, flame and metallic arc site/metals, carbons, fluxes, rods, mask. Guaranteed Fast. Only \$19.95. Used by the Navy. Splendid for fast and MAGIC WELDER MFG. CO., 241 LS Canal fast. New York.



FEWER HANDS ... FASTER HANDLING

Stewedore Jr. Power Belt Conveyor makes produce handling easy and rapid. It takes the grunt and faffgue out of lifting in loading, stacking and stewaring operations. It gets your produce to the consumer faster and in better condition . . . with less manpower expended.

Lightweight and easy to handle, Stevedore Jr. can be moved about to fit changing needs. Fully adisable from 18° level to a delivery height of 64°, it lifts a 25 pound distributed load at the rate of 50° per minute. Stevedore Jr. plugs in on any standard II0 volt lighting circuit to make light work out of handling your boxes, baskets, crates or bags of produce.

Write for detailed information on how you can us faver hands and get faster handling. Don't delay, write today!

The Rapids-Standard Company, Inc. 364 Peoples Natl. Bank Bldg. Grand Rapids 2, Michigan

LITERATURE tells how to

Disinfect with FIRE! **AEROIL Kerosene Torch**

Great for garden spraying, whitewash, cold water paints, disinfectants. Save labor with this tool of 99 uses. Thaw water pipes, split giant rocks. Destroy parasites, green weeds, seeds and all, instantiv. Endorsed by leading Agriculturalists the world over.

600,000 USERS

Omniete with 4 gal. corrusated Tank, Burner and Hose its express collect. 5 gal. \$27.25. Spray attachment \$3.50. Immediate shipment. Descriptive literature FREE.

SINE EQUIPMENT AFG, Quckertown, Pg.

NEW OTTAWA SAW EASY 10 BOVE Polic trees. Sews big long, small togs, Hunbe. Pulley work. Fifty to hand, wright less than lower majes. Fifty to hand, wright less than lower majes. There was the best manners to the first majes. There was the best manners to the first majes. The was mired. CO., \$32 Pine 34., Ottawa, Kanen

BEEKEEPERS' VIEWPOINT

(Continued from page 33)

skilled beekeeper on either a cash or share basis. It seems likely that fruit growers will be more careful about spray poisoning their own bees than they might be of rental bees. In fact, it seems that there is no good reason for having bees in the orchard at any time except during the blooming sea-

Commercial beekeepers are as busy with their schedule as are fruit growers. They have a relatively high investment of from \$15 to \$25 per hive and depend upon a honey crop of \$10 or more per colony, both of which are damaged by spray poisoning. Labor is short. Supplies are hard to get. The bulk of the bees are not located near the fruit belt. The risk of loss from moving the bees to new locations is complicated by the possibility that there may be foulbrood in the orchard area to which the bees are taken, and the bees may not get as much nectar from fruit blooms as they would in the open country from dandelion and other flowers. All in all, the average commercial beekeeper will not be attracted by a low rental price.

Some responsibility must be assumed by fruit growers for loss by spray poisoning. This is a community affair as well as an individual responsibility since bees may fly a mile or more during warm days at blooming time. No matter how careful the fruit grower himself may be, a neighbor who sprays out of schedule may poison all the bees in a square mile.

There are enough draw-backs from the beekeeper's standpoint to warrant his demand for some assurance to cover these unusual risks. It seems that the fruit grower faces the problem of reducing losses from spray poisoning or of financing his own supply of bees for pollination.

STATE NEWS

(Continued from page 20)

The 1946 Proceedings of the Society is now on the press and will soon be in the hands of members, giving them the splendid papers at the Cincinnati meeting and at the

papers at the Cincinnati meeting and at the summer meeting of the Society in 1945. The Ohio fruit industry has applied for membership in the Ohio Council for Farm Cooperatives, and has voted L. G. Dean, Geneva, the voting delegate for fruit growers on the Council. Associated with him as ers on the Council. Associated with him as alternate advisers are H. L. Mantle, Painesville, and Harry W. Lutz, Carroll, for apples, and representing the Ohio Apple Institute; Lester Luchsinger, Port Clinton, representing Ohio peach growers; Ben Davis, Clyde, representing cherry growers; and A. G. Hinz, Avon Lake, representing grape growers. The Council is interested in the overall problems of promoting legislative and marketing programs of best interlative and marketing programs of best interest to both growers and consumers.-F. H. Beach, Sec'y. Ohio State Horticultural Society, Columbus.



BORON in AGRICULTURE

For effective control of BORON deficiency disease, write for a copy of the newly revised "Boron In Agriculture." Pacific Coast Borax Company, 51 Madison Avenue, New York 10, New York.

to them 629 we at all

BORAX-BORIC ACID



HUNT'S GRAFTING WAXES RODENT REPELLENT, ETC. M. J. Beck Co., Box 7, Lansing 1, Mich.

OF A HORTICULTURIST

SPRING—gentle, enchanting, rapturous—brings a period of excitement and stimulated hope. The Vernal Equinox is like the dawn, the break-of-day; it is nature's debut. Then the Universe is more like a maternity ward, for the embryos of new living entities are being born to thousands upon thousands of maternal plants.

Nature in her pregnancy has just given birth to a panorama of flowers and now her myriads of progeny expand, grow and develop. The miniature apples, peaches, grapes and aggregations of tiny drupelets are teeming with activity within their carpel walls. One is to be a Golden Jubilee peach, another a Delicious apple and still another a Premier strawberry, all pre-determined by nature's intricate mechanism of heredity.

The Travail of Spring

The orchardist views the travail of spring just as he would look upon the expectancy of an addition to his own flesh and blood. He is impatient, cautious and alert to every sunshiny day, to every menacing cloud and to every chilling breeze. These he knows too well will influence beyond his control the birth of fruit in his plantings. Nature's cradle is not always the quiet, peaceful and tender hammock the grower would like it to be.

But now most of the vagrancies of the genesis are over and the orchardist can rest a little more at ease. However, he is yet aware that expert care must be given the rudiments of his expected harvest. On every hand pest and pestilence stalk the faultless infants of nature. Every tiny fruit is a "heel of Achilles," and every supporting leaf is no less protected. The orchardist must lay upon them a blanket of spray to protect and shield them from moth, beetle and spore.

Orchardist Protects Trees

He tears away the sod and weeds from around the trees because he knows they will compete for the precious water and nutrients of the soil. A thick bed of mulch he spreads beneath the trees to catch and hold the rainfall and to keep the good earth about the roots moist and cool. A



Appleblossom time in New England.

thirsting apple in its infancy he knows will give up its hold on the tree and fall to the ground, never to attain the rich rewards of maturity.

The growing, expanding "fruitlets" need food and lots of it to stave off starvation and hunger, which if permitted to ensue would result in their abscission and death. Fertilizers, mulches and manure which the orchardist applies to the soil beneath the trees makes more plentiful the nutrients in the soil solution. The hungry rootlets sip of this "rich" solution and gain energy to grow into new and unexploited areas.

Leaves Are Factories

Once inside the root, the nutrient solution diffuses to all parts of the plant, especially to the active green leaves and tiny fruits. The leaves are virtually minute factories, chemical factories in miniature, for here the nutrients of the soil solution, the nitrogen, phosphorus, potassium and many others are worked over into valuable health-giving foods— carbohydrates, fats and proteins. Through its intricate system of conducting tissues the tree transports these foods from the leaves to the hungry, growing fruits, yet in their infancy and youth. As they

grow to maturity they need larger and larger quantities of food to build the walls and protoplasm of new cells and developing tissues. What the fruit doesn't use in growth it stores, and by the time it has reached "old age," it becomes senile and is ready to release its hold upon the tree, it is a rich bundle packed full of stored energy. That is the orchardist's reward for his care and efforts. He knows that the lifeline of water, nutrients and food to the developing fruits must be maintained at all costs and that the little green factories must never experience a dearth of raw materials. He is the vigilant of the night keeping watch over his flock-his orchard, his grove, his vineyard, his berry plantation.

The west

Here lies

Among t

Pacific a

As the

Unpredictable Nature

The orchardist is far from being a sentimentalist, though at times he must cater to nature's whims. But nature sometimes expels her wrath upon him as though beating out vengeance. With subdued emotions, the orchardist must then await patiently the outcome of the "death blow." Perhaps it's this constant experience with unpredictable nature that makes the fruit grower the particular kind of person he is.

SEATTLE MONTANA NO. DAK, MINN. TACOMA WIN CITIES PORTLAND IDAHO OREGON WYÓMING POCATELLO* **OMAHA** OGDEN SALT LAKE EVADA CÓLO. KANS RANCISCO NEW OKLA. ARK MEXICO TEXAS LONG BEACH

Opportunity for Industry

The western territory served by Union Pacific is rich in countless raw materials required by industry; has power and other necessary facilities; has a high type of native-born labor. Here lies the opportunity for industry to build and expand.

Among the principal industrial tracts owned or served by Union Pacific are those located in, or near by, the following cities:

Omaha, Neb.
Council Bluffs, Iowa
St. Joseph, Mo.
Kansas City, Mo.-Kans.
(Fairfax District)
Denver, Colo.
Salt Lake City, Utah

IST

eed larger od to build

f new cells

What the

n it stores, ached "old

is ready to

ree, it is a

of stored

rdist's re-

forts. He

ter, nutri-

oing fruits

costs and

ries must f raw ma-

the night

-his or-

yard, his

n being a times he ms. But er wrath out ventions, the

patiently w." Per-

ence with akes the

kind of

ROWER

Ogden, Utah Portland, Oregon Seattle, Wash. Tacoma, Wash. Spokane, Wash. Los Angeles, Cal. Long Beach, Cal.

As the map shows, all these cities are located on the main lines of the Union Pacific Railroad.


The circles on the map indicate 500-mile distribution areas. In normal times, Union Pacific operates a fast, merchandise freight (L. C. L.) service within these short-haul areas, with door-to-door pick up and delivery service.

These industrial properties are improved and occupied by many diversified manufacturing and commercial enterprises.

Necessary utilities are available and each tract is served by adequate industrial trackage. Whenever required, additional trackage can be provided.

You are cordially invited to write W. H. HULSIZER, General Manager of Properties, Union Pacific Railroad, Omaha 2, Nebraska, regarding industrial districts in territory served by Union Pacific.


be Specific - say "Union Pacific"



The Progressive

UNION PACIFIC RAILROAD
The Strategic Middle Poute

MAY, 1946



A Golden Opportunity

THE SLOGAN—"Food Will Win the War and Write the Peace"—is familiar to us all. At the present time, wheat, flour and other grains are considered the major weapons of peace. The largest export of cereals in our history is planned to help ease the world food crisis that may prove to be the worst in modern times.

Wheat is important because, as Secretary of Agriculture Clinton P. Anderson said recently, "It can be shipped, stored, and rationed to large numbers of people more readily than any other food that will support life and can be obtained in quantity."

Curtailment in the use of flour and bakery products, which is a number one must on the government's program, undoubtedly opens a wider market for fresh commodities such as fruits. In taking the place of cereals, fruit and other fresh commodities can be considered another major weapon of peace.

Fears of surpluses in a country that produces ten per cent of the world's food supply have vanished. There should be a ready market for every fruit grown in 1946.

Trends in Transportation

THE CONSUMPTION of fresh fruits and vegetables is definitely on the upgrade and with it goes the problem of better means of transporting high-quality produce from production areas to consumption centers.

Already improvements are under way in modernizing the present railway refrigerator car through proper refrigerating systems. The Pacific Fruit Express Company, which is jointly owned by the Union Pacific and Southern Pacific railroads, is planning to construct 2,000 new refrigerator cars. In 1945 the refrigerator car committee established by the industry through the United Fresh Fruit & Vegetable Association made a number of suggestions for

new refrigerator cars and the cars being built for the Pacific Fruit Company will largely conform with these recommendations.

Three inches of sidewall ventilation and three and one-half inches at floor and ceiling have been carried in cars constructed during the past year. Insulation details on the new cars have not been definitely announced but it is believed that they will carry four and four and one-half inch insulation, respectively.

Among other features to be included in the new cars are adjustable bunkers, half-stage icing devices, preco air circulating fans and sidewall flues.

The fact that the new cars are to be equipped with steel wheels and easier riding trucks will mean faster schedules for the fresh fruit and vegetable industry and better arrival condition for its produce.

The Open Road

WITH WAR-TIME controls over automobile travel a thing of the past, the fruit grower can again expect his city customers to follow the lure of the open road and stop at his roadside stand for fresh produce.

So it's time to shine up the rusty hinges on your stand, put on a new coat of paint, and brush up on the essentials of roadside marketing.

The first place to start, of course, is in your orchard, vineyard or small fruit planting and grow high-quality fruit. Fresh, high-quality produce has been found to have a far greater drawing power than low prices—so build up a reputation for fruits of distinction.

Your roadside stand needn't be an elaborate affair. Just make sure that it's neat, well-painted, attractively landscaped, and above all, convenient to the consumer. Locate your stand from 30 to 100 feet off the main road so that travelers can drive up to your stand and be out of traffic.

And now that your customers have arrived, how about a pleasing display with plenty of eye appeal to tempt them into buying more than they really came for?

A customer would find it most disconcerting to make a trip to your stand just to find a sign that reads "Closed." So if you don't plan to keep your stand open at all times, be sure to advertise your business hours.

It's important to remember, too, that a pleasing smile and a friendly and courteous word has never handicapped any business. It's a sure-fire way to make customers remember you, your product and your roadside stand.

At Blossom Time

MAYTIME IS blossom time in the orchard. It's a time of intense activity, both obvious and microscopic. It requires about a year and a half to produce a winter variety of apple, for instance. The blossom buds begin to form the latter part of June or early July and may continue longer. They do not open until the following May and the fruit is picked in October. So that the whole process is longer than one usually thinks. But the critical time is when the blossoms reach full bloom.

Now one should not confuse pollination and fertilization. The former is merely the carrying of pollen by insect, wind, or other agency from the stamen to the pistil, either of the same or of another flower, But fertilization is quite a different matter. The pollen grain germinates on the stigma of the pistil and the pollen tube grows down the style to the egg and discharges the male element or sperm. When this sperm or male gamete (marrying cell) unites with the egg, fertilization or sexual propagation occurs. Here we see two nuclei uniting to form one while in other parts of the plant body one cell divides to form two. After fertilization has taken place this first cell is called the "zygote". and it then divides and divides until the embryo in the seed is formed. Thus a new plant is formed from two parents.

But blossom time is a precarious time. So much hangs in the balance. If the temperature is cold or windy the bees and other insects do not fly and pollination does not take place. And unless seeds are produced the blossoms usually fall and no fruit results.

Or, if it is cold enough, the blossoms freeze with a 1945 result. Furthermore, if the temperature is low following pollination, even though it does not freeze, the growth of the pollen tube is so slow that fertilization never occurs.

Then there is the problem of nutrition of the young blossoms. If competition for food supply is great we find the blossoms falling because they starve, and this is apparently a critical matter particularly with certain varieties like Stayman Winesap and Delicious.

So, all told, Nature trembles in the balance about the time you are reading the May issue of the American Fruit Grower. But most of the time "all's right with the world" and a bounteous harvest results. Little wonder fruit growing is an interesting vocation and that unusual people are attracted to it.

n time in the fintense accomicroscopic. It and a half ety of apple, in buds begin of June or tinue longer, the following taked in Occle process is thinks. But the blossoms

confuse poln. The forng of pollen her agency pistil, either ther flower. a different germinates stil and the n the style es the male this sperm rying cell) tilization or s. Here we o form one the plant form two. aken place ne "zygote". ivides until is formed.

precarious he balance. d or windy cts do not s not take re produced and no fruit


rmed from

esult. Furure is low en though wth of the t fertiliza-

lem of nussoms. If ly is great ng because parently a with cern Winesap

membles in the you are the AMERIthe most of the world" at results. ing is an at unusual





It turns the topsoil. It breaks up plow pan. Two basic tillage operations performed at the same time by Oliver's new TNT Plow!

How it does this job is of primary importance to you! Like any other plow, the "TNT" turns the furrow slice in the usual way. But, it loosens the ground underneath the topsoil without bringing sterile subsoil to the surface.

Oliver's specially designed secondary bases break up plow pan and the plant-stunting colloidal silt layer to an adjustable maximum depth of 4 inches. Root penetration is facilitated . . . rainfall run-off is decreased . . . moisture readily sinks into the subsoil where it's stored for dry-spell needs . . . better aeration is provided . . . crop yields are increased.

The OLIVER Corporation

400 West Madison Street, Chicago 6, Illinois



OLIVER



Note how the secondary base breaks up plow pan without disturbing the topsoil.



Ask your Oliver dealer to tell you about the sensational new Oliver TNT Plow. It's a new soil conservation tool that will help save your fertile fields. Learn more about its valuable functions.

"FINEST IN FARM MACHINERY"



Fruit Protection Schedule-

There's more money in the bank each year for growers who-

Growers who wish to ship more fruit and better fruit know that reliable spraying and dusting materials properly applied are the backbone of any fruit protection schedule.

That's why growers stay with Dow products year after year . . . because Dow insecticides are closely checked for uniformity, economy, and ability to do a good job in combination with

each other, which, of course, results in economy.

Together, these products add up to a close-knit, all-season protection plan that you can rely on with confidence. Ask your dealer or state experiment station for complete information.

PRE-BLOSSOM STAGE—Now is the time to aim for the market!

"MIKE" SULFUR—microscopically fine, contains more than 95% active sulfur, wets instantly, covers thoroughly, gives superior finish to fruit, gives extra thorough coverage against apple scab and brown rot. DOW DRY LIME SULFUR is easy to handle, dissolves readily in cold water, imparts finish to fruit. BORDOW is an effective copper fungicide particularly recommended for leaf spot on cherries, prunes and plums. DOW ARSENATE OF LEAD is a reliable spray, controlling most leaf-eating and chewing insects.

Prevent build-up of red mites on apples during the summer months by spraying with *DN*-111. Don't wait till foliage becomes bronzed—start when one or two mites per leaf are present. *DN*-111 gives excellent results either used alone or combined with sulfur or lead arsenate. It acts as a safener for arsenate of lead to reduce foliage injury; and it can be used with App-L-Set for mites present during the preharvest stage. Also available now—Dow DDT-25 Wettable—a concentrated powder for control of codling moth.

